

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS


IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 215/227	A1	(11) International Publication Number: WO 00/34244 (43) International Publication Date: 15 June 2000 (15.06.00)
(21) International Application Number: PCT/US99/28428 (22) International Filing Date: 1 December 1999 (01.12.99) (30) Priority Data: 60/111,079 4 December 1998 (04.12.98) US (71) Applicant: BRISTOL-MYERS SQUIBB COMPANY [US/US]; P.O. Box 4000, Princeton, NJ 08543-4000 (US). (72) Inventors: HEWAWASAM, Piyasena; 31 Brookview Lane, Middletown, CT 06457 (US). STARRETT, John, E., Jr.; 23 Hawks Nest Circle, Middletown, CT 06457 (US). (74) Agent: ALGIERI, Aldo, A.; Bristol-Myers Squibb Company, 5 Research Parkway, P.O. Box 5100, Wallingford, CT 06492-7660 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: 3-SUBSTITUTED-4-ARYLQUINOLIN-2-ONE DERIVATIVES AS POTASSIUM CHANNEL MODULATORS <div style="text-align: center;">  (I) </div> (57) Abstract <p>The present invention provides novel 3-substituted-4-arylquinolin-2-one derivatives having general formula (I) wherein R, R¹, R², R³, R⁴, R⁵, R⁶ and R⁷ are as defined herein, or a nontoxic pharmaceutically acceptable salt thereof which are modulators of the large conductance calcium-activated K⁺ channels and are useful in the treatment of disorders which are responsive to the opening of the potassium channels.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

3-SUBSTITUTED-4-ARYLQUINOLIN-2-ONE
5 DERIVATIVES AS POTASSIUM CHANNEL MODULATORS

FIELD OF THE INVENTION

10

The present invention is directed to novel 3-substituted-4-arylquinolin-2-one derivatives which are modulators of the large-conductance calcium-activated potassium (BK) channels and, therefore, useful in the protection of neuronal cells and diseases arising from
15 dysfunction of cellular membrane polarization and conductance. The present invention also provides a method of treatment with the novel substituted quinolin-2-one derivatives and to pharmaceutical compositions thereof.

20

BACKGROUND OF THE INVENTION

Potassium channels play a key role in regulation of cell
25 membrane potential and modulation of cell excitability. Potassium channels are largely regulated by voltage, cell metabolism, calcium and receptor mediated processes. [Cook, N. S., Trends in Pharmacol. Sciences (1988), 9, 21; and Quast, U., et al., Trends in Pharmacol. Sciences (1989), 10, 431]. Calcium-activated potassium (K_{Ca}) channels
30 are a diverse group of ion channels that share a dependence on intracellular calcium ions for activity. The activity of K_{Ca} channels is regulated by intracellular $[Ca^{2+}]$, membrane potential and

- 2 -

phosphorylation. On the basis of their single-channel conductances in symmetrical K⁺ solutions, K_{Ca} channels are divided into three subclasses: large conductance (BK) > 150 pS; intermediate conductance 50-150 pS; small conductance < 50 pS. Large-conductance calcium-activated potassium (Maxi-K or BK) channels are present in many excitable cells including neurons, cardiac cells and various types of smooth muscle cells. [Singer, J. et al., Pflugers Archiv. (1987) 408, 98; Baro, I., et al., Pflugers Archiv. (1989) 414 (Suppl. 1), S168; and Ahmed, F. et al., Br. J. Pharmacol. (1984) 83, 227].

10

Potassium ions play a dominant role in controlling the resting membrane potential in most excitable cells and maintain the transmembrane voltage near the K⁺ equilibrium potential (E_K) of about -90 mV. It has been shown that opening of potassium channels shift the cell membrane potential towards the equilibrium potassium membrane potential (E_K), resulting in hyperpolarization of the cell. [Cook, N. S., Trends in Pharmacol. Sciences (1988), 9, 21]. Hyperpolarized cells show a reduced response to potentially damaging depolarizing stimuli. BK channels which are regulated by both voltage and intracellular Ca²⁺ act to limit depolarization and calcium entry and may be particularly effective in blocking damaging stimuli. Therefore cell hyperpolarization via opening of BK channels may result in protection of neuronal cells.

A range of synthetic and naturally occurring compounds with BK opening activity have been reported. The avena pyrone extracted from avena sativa-common oats has been identified as a BK channel opener using lipid bi-layer technique [International Patent application WO 93/08800, published May 13, 1993]. 6-Bromo-8-(methylamino)imidazo[1,2-a]pyrazine-2-carbonitrile (SCA-40) has been described as a BK channel opener with very limited electrophysiological experiments [Laurent, F. et al., Br. J. Pharmacol. (1993) 108, 622-626]. The flavanoid,

30

- 3 -

Phloretin has been found to increase the open probability of Ca^{2+} -activated potassium channels in myelinated nerve fibers of Xenopus laevis using outside-out patches [Koh, D-S., et al., Neuroscience Lett. (1994) 165, 167-170].

5

In European patent application EP-477,819 published January 4, 1992 and corresponding U.S. Patent No. 5,200,422, issued April 6, 1993 to Olesen, et al., a number of benzimidazole derivatives were disclosed as openers of BK channels by using single-channel and
10 whole-cell patch-clamp experiments in aortic smooth muscle cells. Further work was reported by Olesen, et al., in European J. Pharmacol., 251, 53-59 (1994).

A number of substituted oxindoles have been disclosed as
15 openers of BK channels by P. Hewawasam, et al., in U.S. Patent No. 5,565,483, issued October 15, 1996.


Sit, et al., in International Patent Application WO 98/23273, published June 4, 1998, and corresponding U.S. Patent No. 5,892,045,
20 issued April 6, 1999, disclosed a series of 4-aryl-3-hydroxyquinolin-2-one derivatives, while Hewawasam, et al., in International Patent Application WO 99/09983, published March 4, 1999, disclosed a series of 4-aryl-3-aminoquinolini-2-one derivatives which are openers of BK channels and useful in the treatment of disorders sensitive to potassium channel
25 opening activity.

E. S. Hamanaka in U.S. Patent No. 5,565,472, issued October 15, 1996, discloses a number of 4-aryl-3-(heteroarylureido)-1,2-dihydro-2-oxo-quinoline derivatives which are inhibitors of acyl coenzyme A;
30 cholesterol acyltransferase and are useful as hypolipidemic and antiatherosclerosis agents.

It is the object of the present invention to provide novel compounds that will modulate potassium channels, in particular, large-conductance calcium-activated potassium (BK) channels which will be useful in diseases arising from dysfunction of cellular membrane polarization and conductance.

SUMMARY OF THE INVENTION


The present invention provides novel 3-substituted-4-arylquinolin-
10 2-one derivatives having the general formula



wherein R, R¹, R², R³, R⁴, R⁵, R⁶ and R⁷ are as defined below, or a non-toxic pharmaceutically acceptable salt thereof which are openers of the large conductance calcium-activated K⁺ channels also known as Maxi-K or BK channels. The present invention also provides pharmaceutical compositions comprising said quinolin-2-one derivatives and to the method of treatment of disorders sensitive to potassium channel opening activity such as ischemia, stroke, convulsions, epilepsy, asthma, irritable bowel syndrome, migraine, traumatic brain injury, spinal cord injury, sexual dysfunction and urinary incontinence.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides novel 3-substituted-4-arylquinolin-2-one derivatives which are potent openers of the high conductance, calcium-activated K⁺-channels (BK channel) and which have the formula



wherein

- 10 R and R¹ each are independently hydrogen or methyl;
 R², R³
 and R⁴ each are independently hydrogen, halogen, nitro or trifluoromethyl, provided R², R³, and R⁴ are not all hydrogen;
- 15 R⁵ is bromo, chloro or nitro;
 R⁶ is hydrogen or fluoro;
 n is an integer from 0 to 6;
 m is an integer of 0 or 1; and
 R⁷ is CH₃, -CRR¹OH, -CHO, -C=NOH, -COCH₃ or aryl
- 20 optionally substituted by one or two substituents selected from the group consisting of halogen, hydroxy, methoxy, amino, acetylamino and trifluoromethyl;
 or a nontoxic pharmaceutically acceptable salt thereof.

25 The present invention also provides a method for the treatment or alleviation of disorders associated with BK channels, such as ischemia, stroke, convulsions, epilepsy, asthma, irritable bowel syndrome,

- 6 -

migraine, traumatic brain injury, spinal cord injury, sexual dysfunction, urinary incontinence and especially male erectile dysfunction which comprises administering together with a conventional adjuvant, carrier or diluent a therapeutically effective amount of a compound of formula I or a
5 nontoxic pharmaceutically acceptable salt thereof.

The term "nontoxic pharmaceutically acceptable salt" as used herein and in the claims is intended to include nontoxic base addition salts with inorganic bases. Suitable inorganic bases such as alkali and
10 alkaline earth metal bases include metallic cations such as sodium, potassium, magnesium, calcium and the like. Unless otherwise specified, the term "halogen" as used herein and in the claims is intended to include bromine, chlorine, iodine and fluorine while the term "halide" is intended to include bromide, chloride and iodide anion.

15

Certain of the compounds of the present invention can exist in unsolvated forms as well as solvated forms including hydrated forms such as monohydrate, dihydrate, hemihydrate, trihydrate, tetrahydrate and the like. The products may be true solvates, while in other cases,
20 the products may merely retain adventitious solvent or be a mixture of solvate plus some adventitious solvent. It should be appreciated by those skilled in the art that solvated forms are equivalent to unsolvated forms and are intended to be encompassed within the scope of the present invention. Certain compounds of Formula I can exist in two
25 tautomeric forms. It should be appreciated by those skilled in the art that when R¹ is hydrogen on the nitrogen atom adjacent to the carbonyl carbon atom, the quinoline ring can exist in an enol form. It is intended that both enolic tautomers of the compounds of Formula I are included within the scope of the present invention.

30

In the method of the present invention, the term "therapeutically effective amount" means the total amount of each active component of

- 7 -

the method that is sufficient to show a meaningful patient benefit, i.e., healing of acute conditions characterized by openers of large conductance calcium-activated K⁺ channels or increase in the rate of healing of such conditions. When applied to an individual active ingredient, administered alone, the term refers to that ingredient alone. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously. The terms "treat, treating, treatment" as used herein and in the claims means preventing or ameliorating diseases, tissue damage and/or symptoms associated with dysfunction of cellular membrane polarization and conductance.

Preferred compounds for use in the method of this invention include the compounds of Formula I listed below:

- 4-(5-chloro-2-methoxyphenyl)-3-(hydroxymethyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
- 4-(5-chloro-2-methoxyphenyl)-3-(hydroxymethyl)-7-(trifluoromethyl)-2(1H)-quinolinone;
- 4-(5-chloro-2-methoxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-quinolinecarboxaldehyde;
- 4-(5-chloro-2-methoxyphenyl)-3-(3-hydroxy-1-propenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
- 4-(5-chloro-2-methoxyphenyl)-3-(3-hydroxypropyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
- 4-(5-chloro-2-hydroxyphenyl)-3-(hydroxymethyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
- 4-(5-chloro-2-hydroxyphenyl)-3-(3-hydroxy-1-propenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
- 4-(5-chloro-2-hydroxyphenyl)-3-(3-hydroxypropyl)-6-(trifluoromethyl)-2(1H)-quinolinone;

- (E)-4-(5-chloro-2-hydroxyphenyl)-3-(2-fluoro-3-hydroxy-1-propenyl)-6-trifluoromethyl)-2(1H)-quinolinone;
- (Z)-4-(5-chloro-2-hydroxyphenyl)-3-(2-fluoro-3-hydroxy-1-propenyl)-6-trifluoromethyl)-2(1H)-quinolinone;
- 5 (E)-4-(5-chloro-2-methoxyphenyl)-3-(2-fluoro-3-hydroxy-1-propenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
- (Z)-4-(5-chloro-2-methoxyphenyl)-3-(2-fluoro-3-hydroxy-1-propenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
- 4-(5-chloro-2-hydroxyphenyl)-3-(2-hydroxyethyl)-6-(trifluoromethyl)-
- 10 2(1H)-quinolinone;
- 4-(5-chloro-2-methoxyphenyl)-3-(2-hydroxyethyl)-6-trifluoromethyl)-2(1H)-quinolinone;
- 4-(5-chloro-2-methoxyphenyl)-3-(4-methoxyphenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
- 15 4-(5-chloro-2-methoxyphenyl)-3-[(4-methoxyphenyl)methyl]-6-(trifluoromethyl)-2(1H)-quinolinone;
- 4-(5-chloro-2-methoxyphenyl)-3-(4-nitrophenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
- 4-(5-chloro-2-methoxyphenyl)-3-(4-aminophenyl)-6-(trifluoromethyl)-
- 20 2(1H)-quinolinone;
- 4-(5-chloro-2-hydroxyphenyl)-3-(3,4-dimethoxyphenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
- 4-(5-chloro-2-hydroxyphenyl)-3-(2,4-dihydroxyphenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
- 25 4-(5-chloro-2-hydroxyphenyl)-3-(4-hydroxyphenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
- 4-(5-chloro-2-hydroxyphenyl)-3-[(4-hydroxyphenyl)methyl]-6-(trifluoromethyl)-2(1H)-quinolinone;
- 4-(5-chloro-2-hydroxyphenyl)-3-(4-acetamidophenyl)-6-(trifluoromethyl)-
- 30 2(1H)-quinolinone;
- 4-(5-chloro-2-hydroxyphenyl)-3-(4-aminophenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;


- 9 -

- 4-(5-chloro-2-hydroxyphenyl)-3-[2-(4-hydroxyphenyl)ethyl]-6-(trifluoromethyl)-2(1H)-quinolinone;
4-(5-chloro-2-hydroxyphenyl)-3-methyl-6-(trifluoromethyl)-2(1H)-quinolinone;
- 5 4-[4-(5-chloro-2-hydroxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)quinolin-3-yl]-3-buten-2-one;
4-(5-chloro-2-hydroxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-quinolinecarboxaldehyde oxime;
4-(5-chloro-2-methoxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-quinolinecarboxaldehyde oxime; and
- 10 4-(5-chloro-2-hydroxyphenyl)-3-(2-hydroxy-2-methylpropyl)-6-(trifluoromethyl)-2(1H)-quinolinone.

The compounds of Formula I may be prepared by various
15 procedures such as those illustrated herein in the examples and in the Reaction Schemes described in the specific embodiments and variations thereof which would be evident to those skilled in the art.

The following Reaction Schemes 1 - 11 illustrate representative
20 general procedures for the preparation of intermediates and methods for the preparation of products according to this invention. It should also be evident to those skilled in the art that appropriate substitution of both the materials and methods disclosed herein will produce the examples illustrated below and those encompassed by the scope of this invention.

25

REACTION SCHEME 1

(a) ClC(O)CH₂CO₂R^a, pyridine, CH₂Cl₂, 0°C to RT


(b) KO^tBu, THF, reflux

(c) Dibal-H, THF-hexanes, -78°C to RT


(d) MnO₂, CH₂Cl₂

The preparation of 2(1H)-quinolinones of Formulae 4 and 5 is illustrated in Reaction Scheme 1. Acylation of the compound of Formula 1 with an acid chloride as shown in the Reaction Scheme provided the amide of Formula 2 wherein R^a is hydrogen or C₁₋₄ alkyl which may be cyclized and dehydrated to the quinolinone of Formula 3 by treatment with a base such as potassium *tert*-butoxide in an inert organic solvent. Exposure of the ester of Formula 3 to a reducing agent such as diisobutyl aluminum hydride affords the primary alcohol of Formula 4 which can then be advantageously oxidized with an oxidant such as manganese dioxide to yield the desired aldehyde of Formula 5.

- 11 -


REACTION SCHEME 2(a) $\text{EtOC(O)CH}_2\text{P(O)(OEt)}_2, \text{NaH, DMF}$ (b) $\text{Dibal-H, THF, } -78^\circ\text{C}$ (c) $\text{PtO}_2, \text{EtOH-HCl, H}_2 (60 \text{ psi})$

As illustrated in Reaction Scheme 2, the homologation of the aldehyde of Formula 5 can readily be accomplished with a phosphonate reagent to produce the unsaturated ester of Formula 6 as a mixture of (E)- and (Z)-isomers which then be separated using column chromatography. Reduction of the ester of Formula 6 may be carried out with a reducing agent such as diisobutyl aluminum hydride to afford the corresponding allylic alcohol of Formula 7. Alternatively, when it is desired to prepare the compound of Formula 9, the ester of Formula 6 is selectively reduced under hydrogenation conditions to reduce the double bond and provide the saturated ester of Formula 8. Treatment of the ester of Formula 8, under conditions similar to the reduction of the ester of Formula 6 will afford the corresponding alcohol of Formula 9.

REACTION SCHEME 3

- (a) BBr_3 , CH_2Cl_2 , -78 to 0°C
 (b) $\text{ClC(O)CH}_2\text{CO}_2\text{Me}$, pyridine, CH_2Cl_2
 (c) KO^tBu , THF, reflux
 (d) Dibal-H, THF, -78°C
 (e) Dibal-H, CH_2Cl_2 , -78°C

In Reaction Scheme 3, the butyloxycarbonyl (BOC) and methyl groups can be removed simultaneously by treatment of the compound of Formula 1a with boron tribromide (BBr_3) to give the aniline of Formula 10. Acylation of the aniline of Formula 10 afforded the corresponding amide of Formula 11, which is readily cyclized and dehydrated under basic conditions with potassium tert-butoxide to provide the lactone of Formula 12. Partial reduction of the lactone with diisobutyl aluminum hydride in THF produced the intermediate lactol of Formula 13. Alternatively, it has been found that by changing solvents from THF to methylene chloride, the lactone of Formula 12 can be reduced with diisobutyl aluminum hydride to provide the desired alcohol of Formula 14.

REACTION SCHEME 4


(a) EtOC(O)CH₂P(O)(OR^a)₂, NaH, DMF

(b) Dibal-H, THF-hexanes, -78°C to RT


(c) PtO₂, EtOH-HCl, H₂ (60 psi)

(d) NaOH, EtOH, RT

- When it is desired to prepare the compound of Formulae 17 and 20, the intermediate lactol of Formula 13 may be treated as shown in
- 5 Reaction Scheme 4 with a phosphonate reagent to afford the unsaturated ester of Formula 15 and then, if desired, be saponified to produce the unsaturated acid of Formula 16. Reduction of the ester of Formula 15 with aluminum hydride affords the corresponding unsaturated alcohol of Formula 17. Alternatively, the hydrogenation of
 - 10 the double bond of the compound of Formula 15 yields the ester of Formula 18, which may be either saponified to produce the acid of Formula 19, or reduced with aluminum hydride to yield the desired alcohol of Formula 20.

REACTION SCHEME 5(a) $\text{EtOC(O)CHFP(O)(OEt)}_2$, NaH, DMF(b) Dibal-H, CH_2Cl_2 , -78°C to RT


- Reaction Scheme 5 illustrates the homologation of the intermediate lactol of Formula **13** with a fluorophosphonate, as described in step (a) in the Reaction Scheme, to afford the unsaturated α -fluoroester ester of Formula **21** as a mixture of (E)- and (Z)-isomers. The crude mixture of esters of Formula **21** may be reduced with aluminum hydride, and the resulting mixture of alcohols are advantageously separated by column chromatography to afford the (E)-olefin of Formula **23** and the (Z)-olefin of Formula **25**. In a similar manner, the aldehyde of Formula **5** which contains a methyl ether can be converted to the corresponding desired olefins of Formulae **24** and **26**.

REACTION SCHEME 6(a) $\text{ClC(O)(CH}_2)_n\text{CO}_2\text{R}^a$, pyridine, CH_2Cl_2 (b) KHMDS, THF, -78°C (c) 35% HBr-AcOH, toluene, $80-90^\circ\text{C}$ (d) pyridine.HCl, $180-200^\circ\text{C}$ (e) $\text{BH}_3\cdot\text{SMe}_2$, THF, -10°C to RT

The preparation of compounds of Formulae **31a** and **31b** is illustrated in Reaction Scheme 6. Acylation of the aniline of Formula **1** with an acid chloride provides the corresponding amide. The amide of Formula **27** may be cyclized under basic conditions to give the dehydrohydroxyquinolinone of Formula **28**, which may then be dehydrated and deesterified under acidic conditions such as HBr/AcOH or pTsOH, to afford the quinolinone of Formula **29**. If desired, removal of the methyl ether can be accomplished with pyridine hydrochloride at elevated temperatures to give the corresponding phenol of Formula **30**. Reduction of the acid of Formula **30** then provides the alcohol of Formula **31a** as the phenol. Alternatively, if the methyl ether of the

phenol is desired, direct reduction of the carboxylic acid of Formula 29 with borane provides the corresponding alcohol of Formula 31b.

REACTION SCHEME 7



(a) $\text{ClC(O)(CH}_2)_n\text{R}^8$, pyridine, CH_2Cl_2

(b) KHMDS, THF, -78°C

(c) H^+


(d) pyridine.HCl, $180\text{--}200^\circ\text{C}$

5

The compound of Formulae 34 and 35 wherein n is 0 to 6 and R⁸ is C₁₋₄ alkyl or aryl optionally substituted by one or two substituents selected from the group consisting of halogen, hydroxy, methoxy, amino, acetyl amino and trifluoromethyl, may be prepared by a similar approach to that outlined in Reaction Scheme 6. Thus, Reaction Scheme 7 illustrates the acylation of the compound of Formula 1, followed by cyclization and dehydration to give the 3-substituted quinoline of Formula 34 as the methyl ether. Demethylation of the compound of Formula 34 with pyridine hydrochloride at elevated temperatures provides the corresponding phenol compound of Formula 35.

10

15


REACTION SCHEME 8

(a) $(\text{MeO})_2\text{P}(\text{O})\text{CH}_2\text{X}$, NaH , DMF


(b) $\text{NH}_2\text{OH} \cdot \text{HCl}$, Et_3N , THF

(c) $\text{NH}_2\text{OH} \cdot \text{HCl}$, an. NaOAc , EtOH

As shown in Reaction Scheme 8, homologation of the intermediate lactol of Formula 13 with a cyano phosphonate or phosphonoacetate reagent provides the corresponding unsaturated nitrile of Formula 36a or acetate of Formula 37a, respectively. Similarly, the methyl ether analogs of Formulae 36b and 37b can be synthesized starting with the aldehyde of Formula 5 and treatment with either a cyano phosphonate or phosphonoacetate reagent, respectively. The oxime of Formula 38a can be prepared from the intermediate lactol of Formula 13 by treating the lactol with hydroxyl amine. Similarly, the methyl ether of Formula 38b, can be prepared from the aldehyde of Formula 5.


REACTION SCHEME 9

- Reaction Scheme 9 illustrates the formation of the lactone of Formula 39 when the hydroxy acid of Formula 30a is treated with a catalytic amount of acid in refluxing toluene. Upon attempted purification of lactone of Formula 39 on silica gel and employing methanol as one of the eluting solvents, the lactone may be converted to the ester of Formula 30b. When it is desired to prepare the substituted alcohol of Formula 40, the lactone of Formula 39 is treated with an excess of a lithium reagent such as methyl lithium to produce the disubstituted alcohol of Formula 40 or alternately, with an equivalent amount to produce a monosubstituted alcohol.

REACTION SCHEME 10

- (a) TIPSCl, Imidazole, DMF
 (b) nBuLi, CH₃I, THF
 (c) TBAF, THF
 (d) K₂CO₃, (CH₃O)₂SO₂, Acetone

The preparation of the N-methyl compounds of Formulae **43** and **44** are depicted in Reaction Scheme 10. Silylation of the alcohol of Formula **31a** with triisopropylsilyl (TIPS) chloride provided the silyl protected ether of Formula **41**. N-alkylation with an alkyl halide such as methyl iodide afforded the compound of Formula **42**, which may be desilylated with a fluoride reagent, step (c), to give the alcohol of Formula **43**. When it is desired to prepare the methylated phenol, the compound of Formula **41** is treated with dimethylsulfate followed by desilylation to afford the dimethyl analog of Formula **44**.

REACTION SCHEME 11

- (a) LiHMDS/THF, -78°C TO RT
 (b) 12N HCl
 (c) pTSA, Toluene, reflux
 (d) LiHMDS
 (e) hv, MeOH


The preparation of the compound of Formula **31a** is advantageously prepared by the reactions illustrated in Reaction

5 Scheme 11. The coumarin compound of Formula **45** is advantageously prepared by condensing γ -butyrolactone with the methyl ester of chlorosalicylic acid which is then readily cyclized with acid to produce the benzopyran-4-one of Formula **46**. Treatment of compound **46** with p-trifluoromethylaniline as illustrated in step (d) produced the

10 dihydrofuran of Formula **47** which is then subjected to photocyclization in an inert organic solvent to afford the compound of Formula **31a**.

- 21 -

In a preferred embodiment of the invention the compounds of Formula I have the formula



- 5 wherein R and R¹ each are independently hydrogen or methyl; R², R³ and R⁴ each are independently hydrogen, halogen, nitro or trifluoromethyl, provided R², R³, and R⁴ are not all hydrogen; R⁵ is bromo, chloro or nitro; R⁶ is hydrogen or fluoro; n is an integer from 0 to 6; m is an integer of 0 or 1; and R⁷ is -CH₃, -CRR¹OH, -CHO, -C=NOH, -COCH₃
- 10 or aryl optionally substituted by one or two substituents selected from the group consisting of halogen, hydroxy, methoxy, amino, acetylamino and trifluoromethyl; or a nontoxic pharmaceutically acceptable salt thereof.
- 15 In another preferred embodiment of the invention, the compounds of the invention include those wherein R and R¹ each are independently hydrogen or methyl; R², R³ and R⁴ each are independently hydrogen, chloro, nitro or trifluoromethyl, provided R², R³ and R⁴ are not all hydrogen; R⁵ is chloro; R⁶ is hydrogen or fluoro; n is 0, 1, or 2; m is 0 or 1;
- 20 and R⁷ is -CH₃, -CH₂OH, -CHO, -C=NOH, -COCH₃ or aryl optionally substituted by halogen, hydroxy, methoxy, amino, acetylamino or trifluoromethyl; or a nontoxic pharmaceutically acceptable salt thereof.

In yet another more preferred embodiment of the invention the

25 compound of Formula I include those wherein R is hydrogen or methyl; R¹ and R⁴ are hydrogen; R² and R³ each are independently

- 22 -

trifluoromethyl; R⁵ is chloro; R⁶ is hydrogen; n is 0, 1, or 2; m is 0 or 1; and R⁷ is -CH₂OH or aryl optionally substituted by halogen, hydroxy, methoxy, amino, acetyl amino or trifluoromethyl; or a nontoxic or a nontoxic pharmaceutically acceptable salt thereof.

5

In another aspect, this invention provides a method for the treatment of or protection from disorders which are mediated by opening of the large conductance calcium-activated K⁺ channels (BK channels) in a mammal in need thereof, which comprises administering to said

10 mammal a therapeutically effective amount of a compound of Formula I or a nontoxic pharmaceutically acceptable salt thereof. Preferably, the compounds of Formula I are useful in the treatment of ischemia, stroke, convulsions, asthma, irritable bowel syndrome, migraine, traumatic brain injury, urinary incontinence, and sexual dysfunction in both men (erectile

15 dysfunction, for example, due to diabetes mellitus, spinal cord injury, radical prostatectomy, psychogenic etiology or any other cause) and women by improving blood flow to the genitalia, especially the corpus cavernosum, and other disorders sensitive to BK channel activating activity.

20

In still another aspect, this invention provides pharmaceutical compositions comprising at least one compound of Formula I in combination with a pharmaceutical adjuvant, carrier or diluent.

25 **Biological Activity**

Potassium (K⁺) channels are structurally and functionally diverse families of K⁺-selective channel proteins which are ubiquitous in cells, indicating their central importance in regulating a number of key cell

30 functions [Rudy, B., Neuroscience, 25: 729-749 (1988)]. While widely distributed as a class, K⁺ channels are differentially distributed as

- 23 -

individual members of this class or as families. [Gehlert, D. R., et al., Neuroscience, 52: 191-205 (1993)]. In general, activation of K⁺ channels in cells, and particularly in excitable cells such as neurons and muscle cells, leads to hyperpolarization of the cell membrane, or in the case of
5 depolarized cells, to repolarization. In addition to acting as an endogenous membrane voltage clamp, K⁺ channels can respond to important cellular events such as changes in the intracellular concentration of ATP or the intracellular concentration of calcium (Ca²⁺). The central role of K⁺ channels in regulating numerous cell functions
10 makes them particularly important targets for therapeutic development. [Cook, N. S., Potassium channels: Structure, classification, function and therapeutic potential. Ellis Horwood, Chichester (1990)]. One class of K⁺ channels, the large-conductance Ca²⁺-activated K⁺ channels (Maxi-K or BK channels), is regulated by transmembrane voltage, intracellular
15 Ca²⁺, and a variety of other factors such as the phosphorylation state of the channel protein. [Latorre, R., et al., Ann. Rev. Physiol., 51: 385-399 (1989)]. The large, single channel-conductance (generally > 150 pS) and high degree of specificity for K⁺ of BK channels indicates that small numbers of channels could profoundly affect membrane conductance
20 and cell excitability. Additionally, the increase in open probability with increasing intracellular Ca²⁺ indicates involvement of BK channels in the modulation of Ca²⁺-dependent phenomena such as secretion and muscular contraction. [Asano, M., et al., J. Pharmacol. Exp. Ther., 267: 1277-1285 (1993)].

25

Openers of BK channels exert their cellular effects by increasing the open probability of these channels [McKay, M. C., et al., J. Neurophysiol., 71: 1873-1882 (1994); and Olesen, S.-P., Exp. Opin. Invest. Drugs, 3: 1181-1188 (1994)]. This increase in the opening of
30 individual BK channels collectively results in the hyperpolarization of cell

- 24 -

membranes, particularly in depolarized cells, produced by significant increases in whole-cell BK-mediated conductance.

The ability of compounds described in the present invention to
5 open BK channels and increase whole-cell outward (K^+) BK-mediated
currents was assessed under voltage-clamp conditions by determining
their ability to increase cloned mammalian (mSlo or hSlo) BK - mediated
outward current heterologously expressed in *Xenopus* oocytes [Butler,
A., et al., Science, 261: 221-224 (1993); and Dworetzky, S. I., et al., Mol.
10 Brain Res., 27: 189-193 (1994)]. The two BK constructs employed
represent nearly structurally identical homologous proteins, and have
proven to be pharmacologically identical in our tests. To isolate BK
current from native (background, non-BK) current, the specific and potent
BK channel-blocking toxin iberitoxin (IBTX) [Galvez, A., et al., J. Biol.
15 Chem., 265: 11083-11090 (1990)] was employed at a supramaximal
concentration (50 nM). The relative contribution of BK channels current
to total outward current was determined by subtraction of the current
remaining in the presence of IBTX (non-BK current) from the current
profiles obtained in all other experimental conditions (control, drug, and
20 wash). It was determined that at the tested concentration the compounds
profiled did not effect non-BK native currents in the oocytes. All
compounds were tested in at least 5 oocytes and are reported at the
single concentration of 20 μ M; the effect of the selected compounds of
Formula I on BK current was expressed as the percent of control IBTX-
25 sensitive current and is listed in Table I. Recordings were accomplished
using standard two-electrode voltage clamp techniques [Stuhmer, W., et
al., Methods in Enzymology, Vol. 207: 319-339 (1992)]; voltage-clamp
protocols consisted of 500-750 ms duration step depolarizations from a
holding potential of -60 mV to +140 mV in 20 mV steps. The
30 experimental media (modified Barth's solution) consisted of (in mM):

NaCl (88), NaHCO₃ (2.4), KCl (1.0), HEPES (10), MgSO₄ (0.82),
Ca(NO₃)₂ (0.33), CaCl₂ (0.41); pH 7.5.

TABLE 1

5

Ex. No.	BK Current*
3	+
4	+
8	++
9	++
17	+
18	+
20	++
21	++
30	++
32	++
34	++
39	++

* at 20 μ M expressed as percent increase over BK current in controls

+ = 100 - 200%

++ = > 200%

10

To determine the ability of these compounds to reduce cell loss resulting from neuronal ischemia, a standard rodent model of permanent focal ischemia, involving occlusion of the middle cerebral artery in the spontaneously hypertensive rat (middle cerebral artery occlusion (MCAO) model) was employed [Tamura, A., et al., Journal of Cerebral Blood Flow and Metabolism, Volume 1, 53-60, (1981)].

15

Selected compounds have been evaluated in the focal stroke model involving permanent MCAO in the spontaneously hypertensive rat. This procedure results in a reliably large neocortical infarct volume that is measured by means of vital dye exclusion in serial slices through the brain 24 hours after MCAO. In the present test, compounds were administered using an intravenous route of administration at 2 hours after occlusion. For example, in this model, the compound of Example 21 reduced the cortical infarct volume by about 25% when administered (0.003 mg/kg) as a single bolus 2 hours after middle cerebral artery occlusion as compared to vehicle-treated (2% DMSO, 98% propylene glycol) control.

The in vivo model on erectile function is described fully in the scientific literature [Rehman, J., Chenven, E., Brink, P. Peterson, B., Wolcott, B., Wen, Y.P., Melman, A., Christ, G.: Diminished neurogenic but not pharmacological erections in the 2- to 3-month experimentally diabetic F-344 rat. Am. J. Physiol. 272: H1960-H1971, (1997)]. Briefly, rats (250-600 g) were anesthetized using sodium pentobarbital, the abdomen opened and the cavernous nerve identified. A pressure catheter was placed in the right corpus cavernosum (crus) to measure intracavernous pressure (ICP). A second catheter was introduced into the carotid artery to measure blood pressure. Test compound (0.1, 0.3 and 1 mg/kg *i.v.*) or vehicle (PEG 400) was given via a catheter placed into the jugular vein.

25

Control intracavernous pressure responses were elicited by electrically stimulating the cavernous nerve via bipolar stimulating electrodes (20 Hz, 0.22 ms pulse width). Stimulus amplitude (0.2 – 20 mA) was adjusted to produce a submaximal intracavernous pressure response (typically 0.2 or 0.5 mA). A series of control intracavernous pressure responses were then obtained using a constant stimulus amplitude. Test compound or vehicle was then administered (200 μ l *i.v.*

30

- 27 -

bolus) and the cavernous nerve was restimulated to evoke a cavernous pressure response at various times post-drug administration. Animals were excluded from the study if the initial ICP responses to nerve stimulation were unstable ("spiky" responses) or if there were time-dependent variations in the magnitude of the control responses. Animals were also excluded if the control ICP/BP response fell outside the 0.3-0.6 range. A repeated measures ANOVA was used for the evaluation of statistical significance.

10 The compound of Example 20 (0.1 – 1 mg/kg) produced an augmentation of the ICP/BP responses elicited by sub-maximal stimulation of the cavernous nerve. A significant increase in the ICP/BP ratio was observed at doses from 0.1-1.0 mg/kg of compound tested.

15 The results of the above biological tests demonstrates that the compounds of the instant invention are potent openers of the large-conductance calcium-activated K⁺ channels (Maxi-K or BK channels). Thus, the compounds of the present invention are useful for the treatment of human disorders arising from dysfunction of cellular
20 membrane polarization and conductance and, preferably, are indicated for the treatment of ischemia, stroke, convulsions, epilepsy, asthma, irritable bowel syndrome, migraine, traumatic brain injury, spinal cord injury, sexual dysfunction, urinary incontinence and especially male
25 erectile dysfunction, other disorders sensitive to BK channel activating activity.

 In another embodiment, this invention includes pharmaceutical compositions comprising at least one compound of Formula I in combination with a pharmaceutical adjuvant, carrier or diluent.

30

 In still another embodiment, this invention relates to a method of treatment or prevention of disorders responsive to opening of potassium

- 28 -

channels in a mammal in need thereof, which comprises administering to said mammal a therapeutically effective amount of a compound of Formula I or a nontoxic pharmaceutically acceptable salt thereof.

- 5 For therapeutic use, the pharmacologically active compounds of Formula I will normally be administered as a pharmaceutical composition comprising as the (or an) essential active ingredient at least one such compound in association with a solid or liquid pharmaceutically acceptable carrier and, optionally, with
10 pharmaceutically acceptable adjuncts and excipients employing standard and conventional techniques.

- The pharmaceutical compositions include suitable dosage forms for oral, parenteral (including subcutaneous, intramuscular, intradermal
15 and intravenous) bronchial or nasal administration. Thus, if a solid carrier is used, the preparation may be tableted, placed in a hard gelatin capsule in powder or pellet form, or in the form of a troche or lozenge. The solid carrier may contain conventional excipients such as binding agents, fillers, tableting lubricants, disintegrants, wetting agents and the
20 like. The tablet may, if desired, be film coated by conventional techniques. If a liquid carrier is employed, the preparation may be in the form of a syrup, emulsion, soft gelatin capsule, sterile vehicle for injection, an aqueous or non-aqueous liquid suspension, or may be a dry product for reconstitution with water or other suitable vehicle before
25 use. Liquid preparations may contain conventional additives such as suspending agents, emulsifying agents, wetting agents, non-aqueous vehicle (including edible oils), preservatives, as well as flavoring and/or coloring agents. For parenteral administration, a vehicle normally will comprise sterile water, at least in large part, although saline solutions,
30 glucose solutions and like may be utilized. Injectable suspensions also may be used, in which case conventional suspending agents may be employed. Conventional preservatives, buffering agents and the like

- 29 -

also may be added to the parenteral dosage forms. Particularly useful is the administration of a compound of Formula I directly in parenteral formulations. The pharmaceutical compositions are prepared by conventional techniques appropriate to the desired preparation
5 containing appropriate amounts of the active ingredient, that is, the compound of Formula I according to the invention. See, for example, Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, PA, 17th edition, 1985.

10 The dosage of the compounds of Formula I to achieve a therapeutic effect will depend not only on such factors as the age, weight and sex of the patient and mode of administration, but also on the degree of potassium channel activating activity desired and the potency of the particular compound being utilized for the particular disorder of
15 disease concerned. It is also contemplated that the treatment and dosage of the particular compound may be administered in unit dosage form and that the unit dosage form would be adjusted accordingly by one skilled in the art to reflect the relative level of activity. The decision as to the particular dosage to be employed (and the number of times to be
20 administered per day) is within the discretion of the physician, and may be varied by titration of the dosage to the particular circumstances of this invention to produce the desired therapeutic effect.

A suitable dose of a compound of Formula I or pharmaceutical
25 composition thereof for a mammal, including man, suffering from, or likely to suffer from any condition as described herein is an amount of active ingredient from about 0.01 $\mu\text{g/kg}$ to 10 mg/kg body weight and preferably, from about 0.1 $\mu\text{g/kg}$ to 5 mg/kg body weight for oral administration. For parenteral administration, the dose may be in the
30 range of 0.1 $\mu\text{g/kg}$ to 1 mg/kg body weight for intravenous administration. The active ingredient will preferably be administered in equal doses from

- 30 -

one to four times a day. However, usually a small dosage is administered, and the dosage is gradually increased until the optimal dosage for the host under treatment is determined.

5 The compounds of the present invention may be employed alone or in combination with other suitable therapeutic agents useful in the treatment of sexual dysfunction such as cGMP PDE inhibitors and particularly cGMP PDE V inhibitors such as sildenafil. Exemplary of the therapeutic agents are PDE V inhibitors selected from imidazo-
10 quinazolines (see WO 98/08848), carbazoles (see WO 97/03675, WO 97/03985 and WO 95/19978), imidazopurinones (see WO 97/19947), benzimidazoles (see WO 97/24334), pyrazoloquinolines (see U.S. Patent 5,488,055), anthranilic acid derivatives (see WO 95/18097), fused heterocycles (see WO 98/07430) and thienopyrimidines (see DE
15 19632423).

 The above therapeutic agents, when employed in combination with the compounds of the present invention, may be used, for example, in those amounts indicated in the Physician's Desk Reference (PDR) or
20 as otherwise determined by one of ordinary skill in the art.

 However, it will be understood that the amount of the compound actually administered will be determined by a physician, in the light of the relevant circumstances including the condition to be treated, the
25 choice of compound of be administered, the chosen route of administration, the age, weight, and response of the individual patient, and the severity of the patient's symptoms.

 The following examples are given by way of illustration and are
30 not to be construed as limiting the invention in any way inasmuch as many variations of the invention are possible within the scope of the invention.

DESCRIPTION OF SPECIFIC EMBODIMENTS

In the following examples, all temperatures are given in degrees Centigrade. Melting points were recorded on a Gallenkamp capillary melting point apparatus are uncorrected. Proton magnetic resonance (^1H NMR) spectra were recorded on a Bruker AC 300 spectrometer. All spectra were determined in the solvents indicated and chemical shifts are reported in δ units downfield from the internal standard tetramethylsilane (TMS) and interproton coupling constants are reported in Hertz (Hz). Splitting patterns are designated as follows: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; br, broad peak; dd, doublet of doublet; bd, broad doublet; dt, doublet of triplet; bs, broad singlet; dq, doublet of quartet. Infrared (IR) spectra using potassium bromide (KBr) were determined on a Perkin Elmer 781 spectrometer from 4000 cm^{-1} to 400 cm^{-1} , calibrated to 1601 cm^{-1} absorption of a polystyrene film and reported in reciprocal centimeters (cm^{-1}). Low resolution mass spectra (MS) and the apparent molecular weight (MH^+) or (M-H^-) was determined on a Finnigan TSQ 7000. The element analyses are reported as percent by weight. Unless otherwise indicated in the Specific Embodiments, R^2 and R^4 are H in the descriptive title of the Examples.

EXAMPLE 1

4-(5-Chloro-2-methoxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-quinolinecarboxylic acid, methyl ester (3, $\text{R}^3 = \text{CF}_3$, $\text{R}^4 = \text{CH}_3$)

Step A: N-[2-[(5-Chloro-2-methoxyphenyl) carbonyl]-4-(trifluoromethyl) phenyl]aminocarboxylic acid, 1,1-dimethylethyl ester

- 32 -

A stirred neat mixture of 4-aminobenzotrifluoride (35 g, 0.218 mol) and (Boc)₂O (52.4g, 0.24 mol) was heated at 80°C for 2 to 3 hours until CO₂ evolution ceased. The mixture was allowed to cool and the ^tBuOH was rotary evaporated. The resultant white solid was recrystallized from
5 hexanes/ether to provide white needles (50.6 g, 89%) of N-(tert-butoxycarbonyl)-4-aminobenzotrifluoride.

Tert-BuLi (130 mL, 0.22 mol, 1.7M in cyclohexane) was added over 20 minutes to a cold (-78°C) stirred solution of N-Boc-4-
10 aminobenzotrifluoride (26.2 g, 0.1 mol) in dry THF (130 mL) under argon. The resultant yellow partial solution was warmed to -45° to -40°C and maintained for 2 hours. The resultant thick yellow slurry of the dianion was cooled to -78°C and neat dry methyl 5-chloro-2-methoxybenzoate (22.1 g, 0.11 mol) was added rapidly. The resultant yellow-brown
15 solution was warmed to -40°C and maintained for 1 hour. The reaction was diluted with ether (200 mL) and quenched with 1N HCl (250 mL) and then allowed to warm to room temperature. The organic layer was separated, washed with water, brine and then dried (Na₂SO₄).
Evaporation of solvents gave a light-yellow solid (49.9 g) which was
20 triturated with ether to afford 31.9 g of the desired titled compound: mp 148-150°C; IR (KBr, cm⁻¹) 3280, 1725, 1640, 1530, 1320, 1250, 1150;

¹H NMR (300 MHz, DMSO-d₆): δ 1.41 (9 H, s), 3.58 (3 H, s), 7.19 (1 H, d, J = 8.9 Hz), 7.49 (1 H, d, J = 2.7 Hz), 7.58 (1 H, d, J = 2.6 Hz), 7.60
25 (1 H, dd, J = 8.9 and 2.7 Hz), 7.93 (1 H, dd, J = 8.7 and 1.9 Hz), 8.12 (1 H, s), 8.15 (1 H, m), 10.35 (1 H, s); MS m/e 430 (MH⁺).

Anal. Calcd. for C₂₀H₁₉ClF₃NO₄: C, 55.88; H, 4.45; N, 3.25.

Found: C, 55.51; H, 4.38; N, 3.26.

- 33 -

Step B: 1-[2-Amino-5-(trifluoromethyl)phenyl]-1'-(5-chloro-2-methoxyphenyl) methanone (1, $R^3 = CF_3$)

To a stirred solution of N-[2-[(5-chloro-2-methoxyphenyl) carbonyl]-4-(trifluoromethyl)phenyl]aminocarboxylic acid, 1,1-dimethylethyl ester (19 g, 0.044 mol) in ethanol (300 mL), 3N HCl was added. The resultant suspension was heated to reflux for 3 hours. The progress of the hydrolysis was monitored by TLC. The reaction mixture was cooled and poured into cold water (500 mL). The product was extracted with ether (2X200 mL) and the combined ether extracts were washed with water, brine and then dried (Na_2SO_4). Evaporation of the ether gave a golden yellow viscous oil which upon standing overnight solidified to afford a beige solid (14.6 g, 100%): mp 90-92°C; IR (KBr, cm^{-1}) 3340, 3470, 1640, 1320, 1240, 1150, 1025;

15

1H NMR (300 MHz, DMSO- d_6): δ 3.68 (3 H, s), 6.97 (1 H, d, $J = 8.8$ Hz), 7.19 (1 H, d, $J = 8.9$ Hz), 7.26 (1 H, d, $J = 1.1$ Hz), 7.36 (1 H, d, $J = 2.7$ Hz), 7.53 (2 H, m), 7.92 (2 H, brd s); MS m/e 330 (MH^+).

20 Anal. Calcd. for $C_{15}H_{11}ClF_3NO_2$: C, 54.64; H, 3.36; N, 4.25.
Found: C, 54.65; H, 3.37; N, 4.16.

Step C: 3-[[2-[(5-Chloro-2-methoxyphenyl)carbonyl]-4-(trifluoromethyl)phenyl]amino]-3-oxopropanoic acid, methyl ester (2, $R^3 = CF_3$, $R^a = CH_3$)

25

A solution of methyl malonyl chloride (1.3 mL, 12 mmol) in anhydrous CH_2Cl_2 (10 mL) was added dropwise to a stirred cold (0°C) solution of 1-[2-amino-5-(trifluoromethyl)phenyl]-1'-(5-chloro-2-methoxyphenyl) methanone prepared in Step B (3.3 g, 10 mmol) and anhydrous pyridine (0.97 mL, 12 mmol) in anhydrous CH_2Cl_2 (30 mL)

30

- 34 -

- under nitrogen. The resultant mixture allowed to warm to room temperature and maintained for 1 hour. The reaction was cooled to 0°C and then quenched with 1N HCl (1 mL). The organic layer was separated and consecutively washed with saturated NaHCO₃, water, 5 brine and then dried (MgSO₄). Evaporation of CH₂Cl₂ gave a beige solid (4.28 g) which was triturated with ether to afford the title compound as a light-yellow solid (3.98 g, 93%); mp 138-140°C; IR (KBr, cm⁻¹) 1120, 1314, 1530, 1644, 1712, 1738;
- 10 ¹H NMR (300 MHz, CDCl₃): δ 3.57 (2 H, s), 3.66 (3 H, s), 3.81 (3 H, s), 6.92 (1 H, d, J = 8.8 Hz), 7.38 (1 H, d, J = 2.6 Hz), 7.47 (1 H, dd, J = 8.8 and 2.6 Hz), 7.65 (1 H, s), 7.75 (1 H, d, J = 8.9 Hz), 8.84 (1 H, d, J = 8.8 Hz), 11.91 (1 H, brd s); MS m/e 428 (M-H)⁻.
- 15 **Step D:** 4-(5-Chloro-2-methoxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-quinolinecarboxylic acid, methyl ester (3, R³ = CF₃, R^a = CH₃)
- Potassium tert-butoxide (0.63 g, 5.6 mmol) was added to a stirred 20 solution of 3-[[2-[(5-chloro-2-methoxyphenyl)carbonyl]-4-(trifluoromethyl)phenyl]amino]-3-oxopropanoic acid, methyl ester prepared in Step C (2.0 g, 4.65 mmol) in anhydrous THF (30 mL) under nitrogen. The resultant mixture was heated to reflux for 30 minutes. The reaction was allowed to cool, diluted with ether (30 mL) and then 25 acidified with 1N HCl (20 mL). The organic layer was separated, washed with brine and then dried (Na₂SO₄). Evaporation of solvents gave a beige solid (1.94 g) which was recrystallized from EtOAc/hexanes to afford the title compound as a white solid (1.82 g, 95%); mp 214-216°C; IR (KBr, cm⁻¹) 1128, 1256, 1322, 1662, 1742;

- 35 -

¹H NMR (300 MHz, CDCl₃): δ 3.68 (3 H, s), 3.70 (3 H, s), 6.97 (1 H, d, J = 8.8 Hz), 7.20 (1 H, d, J = 2.5 Hz), 7.36 (1 H, s), 7.44 (1 H, dd, J = 8.8 and 2.5 Hz), 7.53 (1 H, d, J = 8.6 Hz), 7.73 (1 H, d, J = 8.6 Hz), 12.43 (1 H, brd s); MS m/e 412 (MH⁺).

5

Anal. Calcd. for C₁₉H₁₃ClF₃NO₄: C, 55.42; H, 3.18; N, 3.40.

Found: C, 55.27; H, 2.94; N, 3.30.

EXAMPLE 2

10

4-(5-Chloro-2-methoxyphenyl)-3-(hydroxymethyl)-6-(trifluoromethyl)-2(1H)-quinolinone (4, R³ = CF₃)

A solution of diisobutyl aluminum hydride (2.16 mL of 1M in
15 hexanes, 2.16 mmol) was added dropwise to a cold (-78°C) stirred solution of the compound of Example 1 (0.22 g, 0.54 mmol) in anhydrous THF (10 mL). The mixture was allowed to warm to room temperature and stirred for 2 to 3 hours. Reaction mixture was cooled in an ice-bath and then carefully quenched by dropwise addition of 1N HCl (10 mL).
20 Reaction mixture was diluted with EtOAc (20 mL) and the organic layer was separated, washed with water, brine and then dried (MgSO₄). Evaporation of solvents gave an off-white solid (263 mg) which was triturated with ether to afford the title compound as a white solid (178 mg, 86%); mp 232-235°C; IR (KBr, cm⁻¹) 1126, 1264, 1322, 1654, 3442;

25

¹H NMR (300 MHz, CDCl₃): δ 3.70 (3 H, s), 4.41 (1 H, d, J = 12.5 Hz), 4.53 (1 H, d, J = 12.5 Hz), 7.01 (1 H, d, J = 8.8 Hz), 7.15 (1 H, d, J = 2.6 Hz), 7.33 (1 H, s), 7.47 (1 H, dd, J = 8.8 and 2.6 Hz), 7.52 (1 H, d, J = 8.6 Hz), 7.71 (1 H, d, J = 8.6 Hz), 12.33 (1 H, brd s); MS m/e 384 (MH⁺).

30

- 36 -

Anal. Calcd. for $C_{18}H_{13}ClF_3NO_3$: C, 56.34; H, 3.41; N, 3.65.

Found: C, 55.72; H, 3.44; N, 3.55.

EXAMPLE 3

5

4-(5-Chloro-2-methoxyphenyl)-3-(hydroxymethyl)-7-(trifluoromethyl)-2(1H)-quinolinone (4, $R^3 = R^4 = H$, $R^2 = CF_3$)

Following the general procedure described in Examples 1 and 2,
10 the title compound was prepared.

mp 174-176°C; MS m/e 384 (MH⁺);

¹H NMR (300 MHz, DMSO-d₆): δ 3.64 (3 H, s), 3.88 (1 H, d, J = 11.0 Hz),
15 4.31 (1 H, d, J = 11.0 Hz), 4.70 (1 H, brd s), 7.05 (1 H, d, J = 8.4 Hz), 7.23
(1 H, d, J = 8.9 Hz), 7.29 (1 H, d, J = 2.4 Hz), 7.36 (1 H, d, J = 8.4 Hz),
7.55 (1 H, dd, J = 8.7 and 2.4 Hz), 7.65 (1 H, s), 12.23 (1 H, s).

EXAMPLE 4

20

4-(5-Chloro-2-methoxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-quinolinecarboxaldehyde (5, $R^3 = CF_3$)

Manganese dioxide (0.44 g, 5 mmol) was added to a stirred
25 solution of the compound of Example 2 (384 mg, 1 mmol) in anhydrous
CH₂Cl₂ (10 mL). Resultant suspension was stirred overnight under
nitrogen. Additional MnO₂ (0.44 g, 5 mmol) was added and continued to
stir the suspension until the oxidation is complete (2 to 3 days). The
suspension was filtered through a pad of Celite, washed with additional
30 CH₂Cl₂. Evaporation of the solvent gave the title compound as a light-

- 37 -

yellow solid (206 mg, 54%): mp 238-240°C; IR (KBr, cm⁻¹) 1120, 1268, 1320, 1678, 1707;

¹H NMR (300 MHz, CDCl₃): δ 3.71 (3 H, s), 7.01 (1 H, d, J = 8.9 Hz),
5 7.10 (1 H, d, J = 2.6 Hz), 7.48 (1 H, m), 7.51 (1 H, d, J = 2.6 Hz), 7.59 (1
H, d, J = 8.6 Hz), 7.82 (1 H, dd, J = 8.7 and 1.8 Hz), 10.29 (1 H, s), 12.53
(1 H, brd s); MS m/e 380 (M-H)⁻.

EXAMPLE 5

10

**(E)-3-[4-(5-Chloro-2-methoxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-quinolinyl]-2-propenoic acid, ethyl ester
(6, R³ = CF₃)**

15

To a stirred cold suspension of NaH (84 mg, 2.1 mmol, 60% in mineral oil) in anhydrous DMF (2 mL) a solution of triethyl phosphonoacetate (0.43 g, 1.95 mmol) in DMF (1 mL) was added dropwise under nitrogen. The mixture was stirred for 30 minutes and then neat 4-(5-chloro-2-methoxyphenyl)-1,2-dihydro-2-oxo-6-
20 (trifluoromethyl)-3-quinolinecarboxaldehyde (0.60 g, 1.62 mmol) was added. Resultant mixture was allowed to warm to room temperature and stirred for 3 hours. After this period TLC shows absence of the aldehyde and formation isomeric mixture of the esters. The reaction was cooled in an ice-bath and quenched with 1N HCl. The product was extracted with
25 1:1 ether/EtOAc, washed with saturated NaHCO₃, water, brine and then dried (Na₂SO₄). Evaporation of solvents gave a beige solid (0.765 g) which was recrystallized from EtOAc/hexanes to provide the title compound as a pure trans (E) isomer (0.497 g). Concentration of mother liquor followed by trituration with ether gave additional 123 mg of the
30 ster as a isomeric mixture. Total combined yield of the purified est rs was 0.62 g (86%). Analytical data for the title compound: mp 270-273°C;

- 38 -

IR (KBr, cm^{-1}) 1126, 1284, 1322, 1664, 1713;

^1H NMR (300 MHz, CDCl_3): δ 1.28 (3 H, t, $J = 7.1$ Hz), 3.70 (3 H, s), 4.20 (2 H, q, $J = 7.1$ Hz), 7.04 (1 H, d, $J = 8.9$ Hz), 7.11 (1 H, d, $J = 2.6$ Hz),
5 7.24-7.33 (2 H, m), 7.43 (1 H, s), 7.48-7.52 (2 H, m), 7.76 (1 H, d, $J = 8.6$ Hz), 12.02 (1 H, brd s); MS m/e 450 (M-H) $^-$.

EXAMPLE 6

10 **(E)-4-(5-Chloro-2-methoxyphenyl)-3-(3-hydroxy-1-propenyl)-6-(trifluoromethyl)-2(1H)-quinolinone (7, $\text{R}^3 = \text{CF}_3$)**

To a stirred cold (-78°C) solution of the ester of the compound of Example 5 (0.3 g, 0.66 mmol) in anhydrous THF (9 mL) a Dibal-H
15 solution in hexanes (3 mmol, 3 mL of 1M solution) was added dropwise under nitrogen. The mixture was allowed to warm to room temperature and stirred for 2 hours. The reaction mixture was cooled in ice-bath and then quenched carefully with 1N HCl (10 mL). Ethyl acetate (30 mL) was added, layers separated, washed with water, brine and then dried
20 (MgSO_4). Evaporation of solvents gave a white solid (0.29 g) which was triturated with ether to afford the title compound as an alcohol (234 mg): mp $266-268^\circ\text{C}$;

^1H NMR (300 MHz, $\text{DMSO}-d_6$): δ 3.67 (3 H, s), 3.96 (2 H, m), 4.76 (1 H, t, $J = 5.3$ Hz), 6.11 (1 H, d, $J = 15.7$ Hz), 7.01 (1 H, s), 7.18 (1 H, dt, $J = 15.7$ and 4.6 Hz), 7.27-7.31 (2 H, m), 7.52 (1 H, d, $J = 8.6$ Hz), 7.60 (1 H, dd, $J = 8.9$ and 2.7 Hz), 7.79 (1 H, dd, $J = 8.6$ and 1.8 Hz), 12.36 (1 H, s); MS m/e 408 (M-H) $^-$.

- 39 -

Anal. Calcd. for $C_{20}H_{15}ClF_3NO_3$: C, 58.62; H, 3.69; N, 3.42.

Found: C, 58.50; H, 3.74; N, 3.35.

EXAMPLE 7

5

4-(5-Chloro-2-methoxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-quinolinepropanoic acid, ethyl ester (8, $R^3 = CF_3$)

To a solution of the ester prepared in Example 5 in ethanol and
10 anhydrous HCl in a Parr shaker bottle, PtO_2 (5-10% by weight) was
treated under nitrogen. The resultant suspension was hydrogenated at
60 psi overnight. The catalyst was filtered off and the filtrate was rotary
evaporated to afford the title compound: mp 193-195°C;

15 1H NMR (300 MHz, $CDCl_3$): δ 1.16 (3 H, m), 2.43 (2 H, m), 2.65 (2 H, m),
3.65 (3 H, s), 4.01 (2 H, m), 6.96-7.01 (2 H, m), 7.14 (1 H, s), 7.25 (2 H, s),
7.40-7.43 (2 H, m), 7.60 (1 H, brd s); MS m/e 452 (M-H) $^-$.

EXAMPLE 8

20

4-(5-Chloro-2-methoxyphenyl)-3-(3-hydroxypropyl)-6-(trifluoromethyl)-2(1H)-quinolinone (9, $R^3 = CF_3$)

Following the general procedure described in Example 6, the
25 compound of Example 7 was reduced to provide the title compound.

mp 200-202°C;

1H NMR (300 MHz, $CDCl_3$): δ 1.60 (2 H, m), 2.20 (3 H, brd m), 3.50 (2
30 H, m), 3.63 (3 H, s), 6.97 (1 H, d, $J = 8.8$ Hz), 7.04 (1 H, d, $J = 2.6$ Hz),
7.16 (1 H, s), 7.19 (1 H, s), 7.41 (1 H, dd, $J = 8.8$ and 2.6 Hz), 7.49 (1 H, d,

- 40 -

J = 8.3 Hz), 7.65 (1 H, J = 7.36 Hz), 12.45 (1 H, brd s); MS m/e 412 (MH)⁺.

EXAMPLE 9

5

4-(5-Chloro-2-hydroxyphenyl)-3-(hydroxymethyl)-6-(trifluoromethyl)-2(1H)-quinolinone (14, R³ = CF₃)

Step A: [2-Amino-5-(trifluoromethyl)phenyl](5-chloro-2-hydroxyphenyl)methanone (10, R³ = CF₃)

10

To a cold solution (-78°C) of N-[2-[(5-chloro-2-methoxyphenyl)carbonyl]-4-(trifluoromethyl) phenyl]aminocarboxylic acid, 1,1-dimethylethyl ester prepared in Example 1, Step A (7.0 g, 21.2 mmol) in methylene chloride (60 mL), 1.0 M BBr₃ solution in methylene chloride (46.7 mL, 46.7 mmol) was added dropwise. The resultant red solution was allowed to warm to room temperature and stirred overnight. The reaction mixture was quenched with saturated NaHCO₃ solution. The organic layer was separated and washed with water, brine and then dried (MgSO₄). Evaporation of the solvent gave a yellow-reddish solid which was recrystallized from CH₂Cl₂/hexanes to afford the title compound as a yellow solid (6.58 g, 98%).

15

20

Step B: 3-[[2-[(5-Chloro-2-hydroxyphenyl)carbonyl]-4-(trifluoromethyl)phenyl]amino]-3-oxopropanoic acid, methyl ester (11, R³ = CF₃)

25

To a solution of [2-amino-5-(trifluoromethyl)phenyl](5-chloro-2-hydroxyphenyl)methanone (0.5 g, 1.58 mmol) and pyridin (0.25 mL, 3.17 mmol) in methylene chloride (15 mL), a solution of methyl malonyl chloride (0.34 mL, 3.17 mmol) in methylene chloride (10 mL) was added

30

- 41 -

dropwise at 0°C. The reaction mixture was then allowed to warm to room temperature and stirred for 2 hours. The reaction mixture was acidified with 1N HCl and the organic layer was separated. It was then washed with saturated NaHCO₃ twice, water, brine and dried (MgSO₄).

5 Evaporation of the solvent gave the title compound as a yellowish oil.

Step C: 2-Chloro-6,8-dihydro-11-(trifluoromethyl)-7H-[1]benzopyrano[3,4-c]quinolin-6,7-dione (12, R³ = CF₃)

10 The crude 3-[[2-[(5-chloro-2-hydroxyphenyl)carbonyl]-4-(trifluoromethyl)phenyl]amino]-3-oxopropanoic acid, methyl ester prepared in Step B was dissolved in THF (15 mL) and potassium *t*-butoxide solution (1M in THF, 1.74 mmol, 1.74 mL) was added. The reaction mixture was heated under reflux for 15 minutes. It was then
15 acidified with 1N HCl and the organic layer was separated. The organic layer was washed with water, brine and dried (MgSO₄). Evaporation of the solvent gave a yellow solid which was triturated with ethyl acetate/hexanes to afford the title compound as a yellow solid (0.48 g, 83%); mp >250°C. MS *m/e* 366 (MH⁺).

20

Anal. Calcd. for C₁₇H₇ClF₃NO₃ • 0.5H₂O: C, 54.49; H, 2.15; N, 3.74.

Found: C, 54.10; H, 1.85; N, 3.63.

¹H NMR (DMSO-d₆): δ 7.48 (d, J=8.8 Hz, 1H), 7.55 (d, J=8.5 Hz, 1H),
25 7.76(dd, J=8.9 Hz, 2.2 Hz, 1H), 7.98 (d, J=8.7 Hz, 1H), 8.10 (d, J=2.1 Hz, 1H), 8.42 (s, 1H). IR (KBr, cm⁻¹): 3479, 3074, 1761, 1652, 1630, 1577, 1368, 1325, 1141.

Step D: 4-(5-Chloro-2-hydroxyphenyl)-3-(hydroxymethyl)-6-(trifluoromethyl)-2(1H)-quinolinon (14, R³ = CF₃)
30

- 42 -

To a cold suspension (-78°C) of 2-chloro-6,8-dihydro-11-(trifluoromethyl)-7*H*-[1]benzopyrano[3,4-*c*]quinolin-6,7-dione prepared in Step C (1.0 g, 2.73 mmol) in methylene chloride (20 mL), a solution of Dibal-H (1M in methylene chloride, 13.7 mL, 13.7 mmol) was added dropwise. The reaction mixture was warmed to 0°C and maintained for 3 hours. The reaction mixture was acidified with 1N HCl and extracted with ethyl acetate twice. The organic layer was separated and washed with water, brine and then dried (MgSO₄). Evaporation of the solvent followed by recrystallization of the crude product from ethyl acetate/hexanes afforded the title compound as a white solid (0.7 g, 69%): mp >250°C; MS *m/e* 368 (M-H)⁺.

Anal. Calcd. for C₁₇H₁₁ClF₃NO₃: C, 55.23; H, 3.00; N, 3.79.

Found: C, 56.59; H, 4.02; N, 3.36.

15

¹H NMR (DMSO-*d*₆): δ 3.90 (dd, *J*=10.9 Hz, 5.3 Hz, 1H), 4.36 (dd, *J*=10.9 Hz, 5.6 Hz, 1H), 4.70 (t, *J*=5.4 Hz, 1H), 7.03 (d, *J*=8.7 Hz, 1H), 7.17 (s, 1H), 7.26 (d, *J*=2.7 Hz, 1H), 7.39 (dd, *J*=8.7 Hz, 2.7 Hz, 1H), 7.53 (d, *J*=8.6 Hz, 1H), 7.81 (dd, *J*=8.9 Hz, 1.9 Hz, 1H), 9.95 (s, 1H), 12.31 (s, 1H).

20

EXAMPLE 10

3-[4-(5-Chloro-2-hydroxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-quinolinyl]-2-propenoic acid, ethyl ester (15, R³ = CF₃)

25

Step A: 2-Chloro-6,8-dihydro-6-hydroxy-11-(trifluoromethyl)-7*H*-[1]benzopyrano[3,4-*c*]quinolin-7-one (13, R³ = CF₃)

30

- 43 -

To a cold solution (-78°C) of 2-chloro-6,8-dihydro-11-(trifluoromethyl)-7*H*-[1]benzopyrano[3,4-*c*]quinolin-6,7-dione prepared in Example 9, Step C (1.15 g, 3.15 mmol) in THF (30 mL), a solution of Dibal-H (1M in THF, 15.7 mL, 15.7 mmol) was added dropwise. The reaction mixture was maintained at -78°C for 4 hours. The reaction mixture was acidified with 1N HCl and extracted with ethyl acetate twice. The organic layer was separated and washed with water, brine and then dried (MgSO₄). Evaporation of the solvent followed by trituration of crude product with ethyl acetate to afforded the title compound as a white solid (0.9 g, 78%): mp >260°C; MS *m/e* 366 (M-H)⁻.

Anal. Calcd. For C₁₇H₉ClF₃NO₃ • 0.25H₂O: C, 54.86; H, 2.57; N, 3.76.

Found: C, 54.92; H, 2.92; N, 3.46.

¹H NMR (DMSO-*d*₆): δ 6.40 (d, J=6.2 Hz, 1H), 7.31 (d, J=8.7 Hz, 1H), 7.55-7.63 (m, 3H), 7.94 (d, J=8.7 Hz, 1H), 8.07 (d, J=2.4 Hz, 1H), 8.38 (s, 1H), 12.44 (s, 1H). IR (KBr, cm⁻¹): 3300, 1669, 1631, 1605, 1575, 1326, 1279, 1133.

Step B: 3-[4-(5-Chloro-2-hydroxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-quinoliny]-2-propenoic acid, ethyl ester (15, R³ = CF₃)

To a cold suspension (0°C) of NaH (60% in mineral oil, 41 mg, 1.0 mmol) in DMF (3 mL), triethyl phosphonoacetate (0.1 mL, 0.5 mmol) was added dropwise. The reaction mixture was stirred at 0°C for 0.5 hours and then a solution of 2-chloro-6,8-dihydro-6-hydroxy-11-(trifluoromethyl)-7*H*-[1]benzopyrano[3,4-*c*]quinolin-7-one prepared in Example 10, Step A (0.15 g, 0.41 mmol) in DMF (5 mL) was added. The red reaction mixture was allowed to warm to room temperature and stirred for 4 hours. The reaction mixture was carefully quenched with 1N

- 44 -

HCl solution and was extracted with ethyl acetate. The organic layer was separated and washed with saturated NaHCO_3 , water, brine and dried (MgSO_4). Evaporation of the solvent followed by recrystallization from ethyl acetate/hexanes afforded the title compound as a white solid (127 mg, 72%): mp 262-268°C (dec.); MS m/e 436 (M-H)⁻.

Anal. Calcd. for $\text{C}_{21}\text{H}_{15}\text{ClF}_3\text{NO}_4$: C, 57.61; H, 3.45; N, 3.20.

Found: C, 57.31; H, 3.46; N, 3.15.

¹H NMR (DMSO-d_6): δ 1.18 (t, J=7.1 Hz, 3H), 4.09 (q, J=7.0 Hz, 1H), 7.08 (d, J=8.8 Hz, 1H), 7.16 (d, J=15.7 Hz, 1H), 7.20 (s, 1H), 7.29 (d, J=2.7 Hz, 1H), 7.34 (d, J=15.7 Hz, 1H), 7.48 (dd, J=8.7 Hz, 2.7 Hz, 1H), 7.57 (d, J=8.6 Hz, 1H), 7.90 (dd, J=8.8 Hz, 1.8 Hz, 1H), 10.1 (s, 1H), 12.6 (s, 1H). IR (KBr, cm^{-1}): 3225, 1683, 1662, 1626, 1323, 1301, 1115.

EXAMPLE 11

3-[4-(5-Chloro-2-hydroxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-quinoliny]-2-propenoic acid (16, $\text{R}^3 = \text{CF}_3$)

20

To a solution of the compound of Example 10 (40 mg, 0.09 mmol) in EtOH (2 mL), 10N NaOH (1mL) was added and the mixture was stirred at room temperature overnight. The reaction was acidified with 1N HCl and the precipitated a light yellow solid of title compound was collected (34 mg, 91% yield): mp 258-261°C; MS m/e 408 (M-H)⁻.

Anal. Calcd. for $\text{C}_{19}\text{H}_{11}\text{ClF}_3\text{NO}_4 \cdot 0.5\text{H}_2\text{O}$: C, 52.79; H, 3.15; N, 3.24.

Found: C, 52.93; H, 2.82; N, 3.10.

¹H NMR (DMSO-d_6): δ 7.07-7.13 (m, 2H), 7.18 (s, 1H), 7.25-7.30 (m, 2H), 7.47 (dd, J=8.7 Hz, 2.7 Hz, 1H), 7.57 (d, J=8.7 Hz, 1H), 7.89 (dd,

- 45 -

J=8.7 Hz, 1.7 Hz, 1H), 10.11 (s, 1H), 12.37 (s, br, 1H), 12.57 (s, 1H). IR (KBr, cm^{-1}): 3144, 2996, 1676, 1628, 1323, 1270, 1252, 1130.

EXAMPLE 12

5

4-(5-Chloro-2-hydroxyphenyl)-3-(3-hydroxy-1-propenyl)-6-(trifluoromethyl)-2(1H)-quinolinone (17, $\text{R}^3 = \text{CF}_3$)

To a cold suspension (-78°C) of the compound of Example 10
10 (0.2 g, 0.46 mmol) in methylene chloride (10 mL), a solution of Dibal-H
(1M in methylene chloride, 2.3 mL, 2.3 mmol) was added dropwise. The
reaction mixture was warmed to room temperature and stirred for 4
hours. The reaction mixture was acidified with 1N HCl and extracted
with ethyl acetate twice. The organic layer was separated and washed
15 with water, brine and dried (MgSO_4). Evaporation of the solvent
followed by recrystallization from ethyl acetate/hexanes afforded the title
compound as an off-white solid (0.14 g, 77%): mp $203\text{--}209^\circ\text{C}$ (dec.); MS
m/e 394 (M-H) $^-$.

20 Anal. Calcd. for $\text{C}_{19}\text{H}_{13}\text{ClF}_3\text{NO}_3 \cdot 0.5\text{H}_2\text{O}$: C, 56.38; H, 3.49; N, 3.46.
Found: C, 56.35; H, 3.72; N, 3.29.

^1H NMR (DMSO-d_6): δ 3.97 (dt, J=1.7 Hz, 4.9 Hz, 2H), 4.77 (t, J=5.3 Hz,
1H), 6.16 (dd, J=15.8 Hz, 1.9 Hz, 1H), 7.05 (d, J=8.7 Hz, 1H), 7.10 (s,
25 1H), 7.16 (d, J=2.7 Hz, 1H), 7.21 (dt, J=15.7 Hz, 4.7 Hz, 1H), 7.41 (dd,
J=8.7 Hz, 2.7 Hz, 1H), 7.51 (d, J=8.5 Hz, 1H), 7.78 (dd, J=8.7 Hz, 1.8 Hz,
1H), 9.90 (s, 1H), 12.32 (s, 1H). IR (KBr, cm^{-1}): 3286, 1656, 1641, 1322,
1294, 1169, 1120, 1075.

EXAMPLE 13

4-(5-Chloro-2-hydroxyphenyl)-1,2-dihydro-2-oxo-6-
(trifluoromethyl)-3-quinolinepropanoic acid, ethyl ester (18,
5 $R^3 = CF_3$)

To a solution of the compound of Example 10 (0.4 g, 0.91 mmol)
in ethanol (20 mL), PtO_2 (38 mg) and 3 drops of 1N HCl were added. The
mixture was hydrogenated in a parr apparatus at 50 psi overnight. The
10 catalyst was filtered off by passing through a pad of Celite and washing
with ethanol. The filtrate was evaporated to dryness and the white
residue was flash chromatographed (silica gel, 2:1 ethyl
acetate:hexanes) to afford the title compound as a white solid (0.29 g,
72%); mp 241-245°C (dec.); MS m/e 438 (M-H)⁻.

15

Anal. Calcd. for $C_{21}H_{17}ClF_3NO_4$: C, 57.35; H, 3.90; N, 3.18.

Found: C, 57.27; H, 4.02; N, 2.99.

1H NMR (CD_3OD): δ 1.18(t, J=7.1 Hz, 3H), 2.49-2.55 (m 2H), 2.69-2.75
20 (m, 2H), 4.05 (q, J=7.1 Hz, 2H), 7.02 (d, J=8.7 Hz, 1H), 7.13 (d, J=2.6 Hz,
1H), 7.26 (s, 1H), 7.38 (dd, J=8.7 Hz, 2.6 Hz, 1H), 7.51 (d, J=8.6 Hz, 1H),
7.73 (dd, J=8.6 Hz, 1.7 Hz, 1H). IR (KBr, cm^{-1}): 3353, 1698, 1656, 1626,
1376, 1311, 1270, 1167, 1128, 1074.

25

EXAMPLE 14

4-(5-Chloro-2-hydroxyphenyl)-1,2-dihydro-2-oxo-6-
(trifluoromethyl)-3-quinolinepropanoic acid (19, $R^3 = CF_3$)

30

To a solution of the compound of Example 13 (38 mg, 0.087
mmol) in EtOH (2 mL), 10N NaOH (1 mL) was added and the mixture

- 47 -

was stirred at room temperature overnight. The reaction was acidified with 1N HCl and the precipitated white solid was collected to afford the title compound (30 mg, 84%): mp 255-258°C; MS m/e 410 (M-H)⁺.

5 Anal. Calcd. for C₁₉H₁₃ClF₃NO₄ • 0.5H₂O: C, 54.24; H, 3.35; N, 3.33.

Found: C, 54.10; H, 3.10; N, 3.28.

¹H NMR (DMSO-d₆): δ 2.32-2.37 (m, 2H), 2.47-2.51 (m, 2H), 7.04-7.07 (m, 2H), 7.25 (d, J=2.6 Hz, 1H), 7.40 (dd, J=8.7 Hz, 2.7 Hz, 1H), 7.52 (d, 10 J=8.6 Hz, 1H), 7.79 (dd, J=8.7 Hz, 1.9 Hz, 1H), 9.98 (s, 1H), 12.10 (s, br, 1H), 12.31 (s, 1H). IR (KBr, cm⁻¹): 3283, 3155, 1714, 1626, 1560, 1405, 1275, 1194, 1167, 1132.

EXAMPLE 15

15

4-(5-Chloro-2-hydroxyphenyl)-3-(3-hydroxypropyl)-6-(trifluoromethyl)-2(1H)-quinolinone (20, R³ = CF₃)

To a cold suspension (-78°C) of the compound of Example 13 (0.2 g, 0.45 mmol) in methylene chloride (10 mL), a solution of Dibal-H (1M in methylene chloride, 3.7 mL, 3.7 mmol) was added dropwise. The reaction mixture was warmed to room temperature and stirred for 4 hours. The reaction mixture was acidified with 1N HCl and extracted with ethyl acetate twice. The organic layer was separated and washed 25 with water, brine and dried (MgSO₄). Evaporation of the solvent followed by recrystallization from ethyl acetate/hexanes afforded the title compound as a white solid (145 mg, 80%): mp 257-259°C (dec.); MS m/e 398 (MH⁺).

- 48 -

Anal. Calcd. for $C_{19}H_{15}ClF_3NO_3 \cdot 0.67 \text{ EtOAc}$:

C, 57.00; H, 4.49; N, 3.07.

Found: C, 57.17; H, 4.62; N, 2.88.

5 ^1H NMR (DMSO- d_6): δ 1.5 (m, 2H), 2.3 (m, 2H), 3.25 (m, 2H), 4.35 (m, 1H), 7.02-7.07 (m, 2H), 7.21 (d, $J=2.5$ Hz, 1H), 7.39 (dd, $J=8.7$ Hz, 2.7 Hz, 1H), 7.51 (d, $J=8.5$ Hz, 1H), 7.77 (d, $J=8.5$ Hz, 1H), 9.90 (s, 1H), 12.24 (s, 1H). IR (KBr, cm^{-1}): 3315, 1654, 1624, 1569, 1324, 1273, 1125, 1073.

10

EXAMPLES 16 and 17

(E)- and (Z)-4-(5-Chloro-2-hydroxyphenyl)-3-(2-fluoro-3-hydroxy-1-propenyl)-6-(trifluoromethyl)-2(1H)-quinolinone
15 (23, $R^3 = \text{CF}_3$) and (25, $R^3 = \text{CF}_3$)

Step A: To a cold suspension (0°C) of NaH (60% mineral oil, 68 mg, 1.7 mmol) in DMF (5 mL), triethyl 2-fluoro-2-phosphonoacetate (0.165 mL, 0.82 mmol) was added. The resultant mixture was stirred at
20 0°C for 0.5 hours and then a solution of 2-chloro-6,8-dihydro-6-hydroxy-11-(trifluoromethyl)-7H-[1]benzopyrano[3,4-c]quinolin-7-one prepared in Example 10, Step A (0.25 g, 0.68 mmol) in DMF (5 mL) was added. The red reaction mixture was allowed to warm to room temperature and stirred for 4 hours. The reaction mixture was quenched with 1N HCl and
25 then extracted with ethyl acetate. The organic layer was separated and washed with saturated NaHCO_3 , water, brine and then dried (MgSO_4). Evaporation of the solvent gave a isomeric mixture of esters of formula (21). (E:Z = 1:2.5, 224 mg, 72%) as a colorless oil.

30 Step B: To a cold suspension (-78°C) of the crude esters from Step A (210 mg, 0.46 mmol) in methylene chloride (10 mL), a solution of

- 49 -

Dibal-H (1M in methylene chloride, 3.3 mL, 3.3 mmol) was added dropwise. The reaction mixture was warmed to room temperature and stirred overnight. The reaction mixture was acidified with 1N HCl and extracted with ethyl acetate twice. The organic layer was separated and washed with water, brine and then dried (MgSO₄). The crude isomeric alcohols were purified by column chromatography (silica gel, 2:1 ethyl acetate/hexanes) to afford individual E-isomer (E)-4-(5-chloro-2-hydroxyphenyl)-3-(2-fluoro-3-hydroxy-1-propenyl)-6-(trifluoromethyl)-2(1H)-quinolinone Example 16 (23, R³ = CF₃) and Z-isomer (Z)-4-(5-chloro-2-hydroxyphenyl)-3-(2-fluoro-3-hydroxy-1-propenyl)-6-(trifluoromethyl)-2(1H)-quinolinone Example 17 (25, R³ = CF₃). The physical characteristics of the (E)- and (Z)-isomer compounds are described below.

Example 16: (E)-4-(5-Chloro-2-hydroxyphenyl)-3-(2-fluoro-3-hydroxy-1-propenyl)-6-(trifluoromethyl)-2(1H)-quinolinone (23, R³ = CF₃)

mp 215-218°C (dec.); MS m/e: 412 (M-H)⁺.

Anal. Calcd. for C₁₉H₁₂ClF₃NO₃ • 0.33H₂O: C, 54.37; H, 3.04; N, 3.34.

Found: C, 54.72; H, 3.11; N, 3.18.

¹H NMR (DMSO-d₆): δ 3.64 (m, 1H), 3.83 (m, 1H), 4.99 (m, 1H), 5.61 (d, J=19.6 Hz, 1H), 7.03 (d, J=8.7 Hz, 1H), 7.20-7.23 (m, 2H), 7.38 (dd, J=8.7 Hz, 2.7 Hz, 1H), 7.54 (d, J=8.7 Hz, 1H), 7.84 (d, J=8.9 Hz, 1H), 10.09 (s, 1H), 12.42 (s, 1H).

- 50 -

Example 17: (Z)-4-(5-Chloro-2-hydroxyphenyl)-3-(2-fluoro-3-hydroxy-1-propenyl)-6-(trifluoromethyl)-2(1H)-quinolinone (25, $R^3 = CF_3$)

5 mp 242-245°C (dec.); MS m/e 412 (M-H)⁻.

Anal. Calcd. for $C_{19}H_{12}ClF_3NO_3 \cdot 0.33H_2O$: C, 54.37; H, 3.04; N, 3.34.

Found: C, 54.62; H, 3.27; N, 3.11.

10 1H NMR (DMSO- d_6): δ 3.81-3.86 (m, 2H), 5.35 (t, $J=5.9$ Hz, 1H), 5.57 (d, $J=40.3$ Hz, 1H), 7.00 (d, $J=8.7$ Hz, 1H), 7.12 (d, $J=2.6$ Hz, 1H), 7.21 (s, 1H), 7.34 (dd, $J=8.7$ Hz, 2.7 Hz, 1H), 7.51 (d, $J=8.6$ Hz, 1H), 7.81 (dd, $J=8.6$ Hz, 1.6 Hz, 1H), 9.90 (s, 1H), 12.32 (s, 1H).

15 **EXAMPLES 18 and 19**

Following the general procedures described for the compounds of Examples 16 and 17, the compounds of Example 18 (24, $R^3 = CF_3$) (E-isomer) and Example 19 (26, $R^3 = CF_3$) (Z-isomer) were prepared from
20 the compound of formula (5) prepared in Example 4, as illustrated in Reaction Scheme 5. The physical characterizing data of the (E)- and (Z)-isomers are described below.

Example 18: (E)-4-(5-Chloro-2-methoxyphenyl)-3-(2-fluoro-3-hydroxy-1-propenyl)-6-(trifluoromethyl)-
25 2(1H)-quinolinone (24, $R^3 = CF_3$)

mp 180-182°C; MS m/e: 426 (M-H)⁻.

30 1H NMR ($CDCl_3$): δ 3.68 (3 H, s), 4.10 (2 H, d, $J = 24.0$ Hz), 5.54 (1 H, d, $J = 18.0$ Hz), 7.01 (1 H, d, $J = 8.9$ Hz), 7.04 (1 H, d, $J = 2.4$ Hz), 7.32 (1 H,

- 51 -

s), 7.46 (1 H, dd, J = 8.9 and 2.3 Hz), 7.52 (1 H, d, J = 8.5 Hz), 7.73 (1 H, d, J = 7.6 Hz), 11.65 (1 H, brd s).

Example 19: (Z)-4-(5-Chloro-2-methoxyphenyl)-3-(2-fluoro-3-hydroxy-1-propenyl)-6-(trifluoromethyl)-2(1H)-quinolinone (26, R³ = CF₃)

mp 230-232°C; MS m/e: 426 (M-H)⁻.

¹H NMR (CDCl₃): δ 3.69 (3 H, s), 4.09 (2 H, d, J = 10.7 Hz), 5.70 (1 H, d, J = 38.1 Hz), 6.98 (1 H, d, J = 8.8 Hz), 7.09 (1 H, s), 7.35 (1 H, s), 7.41 (1 H, d, J = 8.8 Hz), 7.50 (1 H, m), 7.67 (1 H, brd s), 11.75 (1 H, brd s).

EXAMPLE 20

4-(5-Chloro-2-hydroxyphenyl)-3-(2-hydroxyethyl)-6-(trifluoromethyl)-2(1H)-quinolinone (31a, R³ = CF₃, n = 2)

Step A: 4-[[2-[(5-Chloro-2-methoxyphenyl)carbonyl]-4-(trifluoromethyl)phenyl]amino]-4-oxobutanoic acid, methyl ester (27, R³ = CF₃, R^a = CH₃, n = 2)

Neat 3-carbomethoxypropionyl chloride (4.8 mL, 0.039 mol) was added to a stirred cold (0°C) solution of aminobenzophenone

1-[2-amino-5-(trifluoromethyl)phenyl]-1'-(5-chloro-2-methoxyphenyl) methanone prepared in Example 1, Step B (7.0 g, 0.021 mol) and anhydrous pyridine (4.8 mL, 0.059 mol) in anhydrous CH₂Cl₂ (80 mL).

The resultant mixture was allowed to warm to room temperature and maintained for 12 hours. The reaction was acidified with 1N HCl (50

mL). The organic layer was separated and washed consecutively with saturated NaHCO₃, water, brine and then dried (MgSO₄). Evaporation of

- 52 -

the CH₂Cl₂ and trituration of the resulting residue afforded 7.71 g (82%) of amide of the title compound.

¹H NMR (300 MHz, CDCl₃): δ 2.7 (4 H, m), 3.06 (3H, s), 3.63 (3H, s),
5 6.85 (1H, d, J = 8.9 Hz), 7.16 (1H, s), 7.30 (1H, d, J = 2.6 Hz), 7.39 (1H,
dd, J = 8.8, 2.6 Hz), 7.57 (1H, s), 7.66 (1H, dd, J = 8.9, 1.9 Hz), 8.79 (1H,
d, J = 8.8 Hz); MS m/e 444 (MH⁺)

Step B: 4-(5-Chloro-2-methoxyphenyl)-4-hydroxy-1,2,3,4-
10 tetrahydro-2-oxo-6-(trifluoromethyl)-3-quinolineacetic acid,
methyl ester (28, R³ = CF₃, R^a = CH₃, n = 2)

A solution of potassium bis(trimethylsilyl)amide (0.5 M in toluene,
57 mL, 28.5 mmol) was added to a stirred cold (-78°C) solution of 4-[[2-
15 [(5-chloro-2-methoxyphenyl)carbonyl]-4-(trifluoromethyl)phenyl]amino]-
4-oxobutanoic acid, methyl ester prepared in Step A (4.05 g, 9.1 mmol)
in anhydrous THF (25 mL) and maintained at -78°C for 3 hours. Acidic
work-up with 1N HCl and followed by extraction with EtOAc afforded the
crude title compound (4.05 g, 100%).

20

Step C: 4-(5-Chloro-2-methoxyphenyl)-1,2-dihydro-2-oxo-6-
(trifluoromethyl)-3-quinolineacetic acid (29, R³ = CF₃, n = 2)

A stirred suspension of crude 4-(5-chloro-2-methoxyphenyl)-4-
25 hydroxy-1,2,3,4-tetrahydro-2-oxo-6-(trifluoromethyl)-3-quinolineacetic
acid, methyl ester prepared in Step B (2 g, 4.5 mmol) in toluene (25 mL)
was treated with a solution of 35% HBr in acetic acid (5 mL). The
resultant mixture was heated at 85°C overnight. The reaction mixture
was evaporated to dryness and the residue was partitioned between

water and EtOAc. The EtOAc extract was washed with brine and dried (MgSO₄) and then evaporated to afford the title compound (1.45 g, 78%).

Step D: 4-(5-Chloro-2-hydroxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-quinolineacetic acid (30, R³ = CF₃, n = 2)

A neat mixture of crude 4-(5-chloro-2-methoxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-quinolineacetic acid prepared in Step C (1.45 g, 3.5 mmol) and pyridinium hydrochloride (5 g, 43.3 mmol) was heated at 185°C for 3 hours. Reaction mixture was allowed to cool and 1N HCl was added and then extracted with EtOAc to afford the title compound (1.20 g, 86%): mp 158-160°C;

¹H NMR (300 MHz, CD₃OD): δ 3.21 (1H, d, J = 16.7 Hz), 3.60 (1H, d, J = 16.7 Hz), 7.0 (1H, d, J = 8.8 Hz), 7.13 (1H, d, J = 2.6 Hz), 7.31 (1H, m), 7.37 (1H, dd, J = 8.8, 2.6 Hz), 7.52 (1H, d, J = 8.6 Hz), 7.75 (1H, d, J = 8.6 Hz); MS m/e 398 (MH⁺).

Step E: 4-(5-Chloro-2-hydroxyphenyl)-3-(2-hydroxyethyl)-6-(trifluoromethyl)-2(1H)-quinolinone (31a, R³ = CF₃, n = 2)

A solution of borane-methyl sulfide complex (2M in THF, 125 mL, 0.25 mol) was added dropwise over 20 minutes to a stirred cold (-10°C) partial solution of the acid 4-(5-chloro-2-hydroxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-quinolineacetic acid prepared in Step D (20 g, 0.05 mol) in anhydrous THF (125 mL) under nitrogen. The resultant clear reaction mixture was allowed to warm to room temperature and continued to stir for 2 to 3 days (HPLC analysis shows absence of starting material). The reaction mixture was cooled in an ice-bath and then quenched with dropwise addition of 1N NaOH (125 mL) until basic and then acidified with 1N HCl. Ether (250 mL) was added and layers

- 54 -

separated, washed with water, brine and then dried (Na_2SO_4).

Evaporation of the solvents gave a brown solid (21.4 g) which was recrystallized from EtOAc-MeOH to afford 2.6 g of pure white solid as the first crop. Trituration of concentrated mother liquor with ether afforded

- 5 8.7 g of off-white solid as the second crop. The second crop was recrystallized from EtOAc-MeOH and combined with first crop to afford a total of 11.1 g of the title compound as a white solid: mp 255-256°C;

- ^1H NMR (300 MHz, CD_3OD): δ 2.73 (2H, m), 3.64 (2H, t, $J = 7.4$ Hz), 7.0
10 (1H, d, $J = 8.8$ Hz), 7.15 (1H, d, $J = 2.6$ Hz), 7.23 (1H, broad s), 7.36 (1H, dd, $J = 8.8, 2.6$ Hz), 7.48 (1H, d, $J = 8.6$ Hz), 7.71 (1H, dd, $J = 8.6, 1.8$ Hz); MS m/e 384 (MH^+).

- Anal. Calcd. for $\text{C}_{18}\text{H}_{13}\text{ClF}_3\text{NO}_3$: C, 56.34; H, 3.41; N, 3.65.
15 Found: C, 56.18; H, 3.58; N, 3.48.

EXAMPLE 21

- 4-(5-Chloro-2-methoxyphenyl)-3-(2-hydroxyethyl)-6-**
20 **trifluoromethyl-2(1H)-quinolinone (31b, $\text{R}^3 = \text{CF}_3$, $n = 2$)**

The compound of Example 20, Step C was reacted by the general procedure described in Example 20, Step E to afford the title compound.

- 25 mp 219-221°C;

^1H (300 MHz, CDCl_3): δ 7.71 (1H, d, $J = 8.4\text{Hz}$), 7.53-7.46 (m, 2H), 7.23
(1H, s), 7.11 (d, 1H, $J = 2.7$ Hz), 7.02 (d, 1H, $J = 8.7$ Hz), 3.79 (m, 2H),
3.70 (s, 3H), 3.79 (t, 2H); MS m/e 397 (MH^+).

30

- 55 -

Anal. Calcd. for $C_{19}H_{15}ClF_3NO_3$: C, 57.37, H, 3.80, N, 3.52.

Found: C, 57.31, H, 3.94, N, 3.38.

EXAMPLES 22 - 33

5

General procedure for the preparation of compounds of Formulas (34) and (35)

10 **Step A:** Acylation of aminobenzophenone of formula (1).

10

Neat acyl chloride (1.2 equivalent) was added to a stirred cold (0°C) solution of aminobenzophenone of formula (1) (1 equivalent) and anhydrous pyridine (1.3 equivalent) in anhydrous CH_2Cl_2 . The resultant mixture was allowed to warm to room temperature and maintained for 2 to 3 hours. The reaction was acidified with 1N HCl, layers separated and the organic layer washed with saturated NaHCO_3 , water, brine and then dried (MgSO_4). Evaporation of the CH_2Cl_2 afforded the corresponding amide of general formula (32).

15

20 **Step B:** Cyclization of the amide of formula (32).

A solution of potassium bis(trimethylsilyl)amide (0.5 M in toluene, 3 equivalent) was added to a stirred cold (-78°C) solution of the amide of formula (32) (1 equivalent) in anhydrous THF and maintained at -78°C for 3 hours. Acidic work-up with 1N HCl and followed by extraction with EtOAc afforded the quinoline of formula (33).

25

Step C: Dehydration of the quinoline of formula (33).

30

A stirred suspension of crude compound of the formula (33) in toluene was treated with a solution of 35% HBr in acetic acid. The

- 56 -

resultant mixture was heated at 85°C overnight. The reaction mixture was evaporated to dryness and the residue was partitioned between water and EtOAc. The EtOAc extract was washed with brine and dried (MgSO₄) and then evaporated to afford the quinoline of formula (34).

5

Step D: Demethylation of the compound of formula (34).

A neat mixture of crude quinoline of formula (34) (1 equivalent) and pyridinium hydrochloride (5 equivalent) was heated at 185°C for 3 hours. The reaction mixture was allowed to cool and 1N HCl was added and then extracted with EtOAc to afford the corresponding hydroxy compound of formula (35).

10

EXAMPLE 22

15

4-(5-Chloro-2-methoxyphenyl)-3-(4-methoxyphenyl)-6-trifluoromethyl)-2(1H)-quinolinone (34a, R³ = CF₃, R⁶ = 4-methoxyphenyl, n = 0)

20 mp 130-135°C;

¹H NMR (300 MHz, CDCl₃): δ 7.62 (1H, d, J = 8.7 Hz), 7.36-7.25 (3H, m), 7.10 (2H, d, J=8.7 Hz), 6.93(1H, d, J = 2.7 Hz), 6.81(1H, d, J = 8.7Hz), 6.77(2H, d, J = 8.7 Hz), 3.77(s, 3H), 3.62(s, 3H) ; MS m/e 459 (MH⁺).

25

Anal. Calcd. for C₂₄H₁₇ClF₃NO₃: C, 62.69, H, 3.73, N, 3.05.

Found: C, 62.74, H, 3.92, N, 2.89.

EXAMPLE 23

4-(5-Chloro-2-methoxyphenyl)-3-[(4-methoxyphenyl)methyl]-
6-(trifluoromethyl)-2(1H)-quinolinone (34b, $R^3 = CF_3$, $R^8 =$
5 4-methoxyphenyl, $n = 1$)

mp 110-114°C;

1H NMR (300 MHz, $CDCl_3$): δ 7.62 (1H, d, $J = 8.7$ Hz), 7.47-7.43 (1H, dd,
10 $J = 2.7$ Hz and 8.7 Hz), 7.29 (d, 1H, $J = 2.7$ Hz), 7.23 (s, 1H), 7.00-6.92
(m, 3H), 6.70 (d, 1H, $J = 8.7$ Hz), 3.78 (dd, 2H), 3.72 (s, 3H), 3.57 (s, 3H);
MS m/e 473 (MH^+).

EXAMPLE 24

15

4-(5-Chloro-2-methoxyphenyl)-3-(4-nitrophenyl)-6-
(trifluoromethyl)-2(1H)-quinolinone (34c, $R^3 = CF_3$, $R^8 =$
4-nitrophenyl, $n = 0$)

20 mp 218-22°C;

1H (300 MHz, $CDCl_3$): δ 8.12 (2H, d, $J = 8.7$ Hz), 7.72 (1H, d, $J = 8.7$ Hz),
7.43-7.36(m, 4H), 7.33-7.29(dd, 1H, $J = 2.7$ and 8.7 Hz), 6.95 (d, 1H, $J =$
2.7 Hz), 6.82 (d, 1H, $J = 8.7$ Hz); MS m/e 474 (MH^+).

25

Anal. Calcd. for $C_{23}H_{14}ClF_3N_2O_4$: C, 58.18, H, 2.97, N, 5.90.

Found: C, 57.70, H, 3.20, N, 5.65.

- 58 -

EXAMPLE 25

4-(5-Chloro-2-methoxyphenyl)-3-(4-aminophenyl)-6-
(trifluoromethyl)-2(1H)-quinolinone (34d, $R^3 = CF_3$, $R^8 =$
5 4-aminophenyl, $n = 0$)

mp 287°C;

1H (300 MHz, $CDCl_3$): δ 7.62 (1H, d, $J = 8.4$ Hz), 7.44 (1H, d, $J = 8.4$ Hz),
10 7.31 (s, 1H), 7.27-7.23 (m, 4H), 6.94-6.79 (dd, 1H, $J = 3.6$ and 8.7 Hz),
6.82 (s, 1H), 6.52 (d, 1H, $J = 8.7$ Hz); MS m/e 444 (MH^+).

Anal. Calcd. for $C_{23}H_{16}ClF_3NO_2$: C, 62.10, H, 3.63, N, 6.30.

Found: C, 61.89, H, 3.81, N, 6.06.

15

EXAMPLE 26

4-(5-Chloro-2-methoxyphenyl)-3-methyl-6-trifluoromethyl-
2(1H)-quinolinone (34e, $n = 0$, $R^8 = Me$, $R^3 = CF_3$)
20

A solution of the corresponding compound of formula (33) (5.63
mmol), 33% HBr in AcOH (38.3 mmol) and 10 mL of AcOH was heated to
75°C for 3 hours. The solution was cooled to room temperature and
quenched with H_2O (50 mL) and then agitated for 12 hours. The
25 precipitate was filtered, washed with H_2O , and dried *in vacuo*. The pale
brown solid was recrystallized from ethyl acetate/hexane to give the title
compound as a white solid (0.550 g, 27% yield).

1H NMR (300 MHz, $CDCl_3$): δ 2.04 (s, 3H), 3.72 (s, 3H), 7.03 (d, 1H, $J =$
30 9.0 Hz), 7.12 (s, 1H), 7.44 (m, 3H), 7.65 (d, 1H, $J = 8.4$ Hz), 10.93 (br s,
1H); MS m/e 368 (MH^+);

- 59 -

Anal. Calcd. for $C_{18}H_{13}ClF_3NO_2 \cdot 0.33 H_2O$: C, 58.79; H, 3.56; N, 3.81.

Found: C, 58.89; H, 3.82; N, 3.53.

EXAMPLE 27

5

4-(5-Chloro-2-hydroxyphenyl)-3-(3,4-dimethoxyphenyl)-6-(trifluoromethyl)-2(1H)-quinolinone (35a, $R^3 = CF_3$, $R^8 = 3,4$ -dimethoxyphenyl, $n = 0$)

10 mp 140-142°C;

1H NMR (300 MHz, $CDCl_3$): δ 7.65 (1H, d, $J = 8.7$ Hz), 7.37 (1H, d, $J = 8.7$ Hz), 7.35 (1H, s), 7.29-7.25 (1H, dd, $J = 2.7$ Hz and $J = 8.7$ Hz), 6.96 (1H, d, $J = 2.4$ Hz), 6.87-6.76 (2H, m), 6.63 (1H, d, $J = 1.8$ Hz), 3.85 (s, 3H), 3.69 (s, 3H), 3.62 (3H, s); MS m/e 489 (MH^+).

15

Anal. Calcd. for $C_{25}H_{19}ClF_3NO_4$: C, 61.30, H, 3.91, N, 2.86.

Found: C, 61.42, H, 3.89, N, 2.75.

20

EXAMPLE 28

4-(5-Chloro-2-hydroxyphenyl)-3-(2,4-dihydroxyphenyl)-6-(trifluoromethyl)-2(1H)-quinolinone (35b, $R^3 = CF_3$, $R^8 = 2,4$ -dihydroxyphenyl, $n = 0$)

25

mp 295°C(dec.);

1H NMR (300 MHz, CD_3OD): δ 7.73 (1H, d, $J = 8.7$ Hz), 7.49 (1H, d, $J = 8.7$ Hz), 7.31-7.27 (2H, dd, $J = 2.7$ and $J = 8.7$ Hz), 7.03-7.00 (2H, m), 6.89 (1H, d, $J = 8.7$ Hz), 6.53-6.49 (2H, m); MS m/e 462 (MH^+).

30

- 60 -

Anal. Calcd. for $C_{22}H_{13}ClF_3NO_4$: C, 59.01, H, 2.93, N, 3.13.

Found: C, 58.38, H, 3.15, N, 2.96.

EXAMPLE 29

5

4-(5-Chloro-2-hydroxyphenyl)-3-(4-hydroxyphenyl)-6-(trifluoromethyl)-2(1H)-quinolinone (35c, $R^3 = CF_3$, $R^8 = 4$ -hydroxyphenyl, $n = 0$)

10 mp 220-240°C;

1H NMR (300 MHz, $CDCl_3$): δ 7.53 (1H, d, $J = 8.7$ Hz), 7.20 (1H, d, $J = 8.7$ Hz), 7.24-7.17 (2H, m), 6.97-6.88 (4H, dd, $J = 8.7$ and 1.9 Hz), 6.81 (1H, d, $J = 8.7$ Hz), 6.77 (2H, d, $J = 8.7$ Hz), 3.77 (s, 3H), 3.62 (s, 3H);

15 MS m/e 459 (MH^+).

EXAMPLE 30

4-(5-Chloro-2-hydroxyphenyl)-3-[(4-hydroxyphenyl)methyl]-6-(trifluoromethyl)-2(1H)-quinolinone (35d, $R^3 = CF_3$, $R^8 = 4$ -hydroxyphenyl, $n = 1$)

20

mp 242-250°C;

25 1H NMR (300 MHz, CD_3OD): δ 7.73 (1H, d, $J = 8.7$ Hz), 7.70 (1H, d, $J = 8.7$ Hz), 7.50 (d, 1H, $J = 8.7$ Hz), 7.42-7.27 (m, 2H), 6.70-6.87 (m, 2H), 6.82 (d, 2H, $J = 8.7$ Hz), 6.56 (d, 2H, $J = 8.1$ Hz), 3.91-3.51 (2H, dd, $J = 13.8$ and 14.7 Hz); MS m/e 445 (MH^+).

30

- 61 -

Anal. Calcd. for $C_{23}H_{15}ClF_3NO_3 \cdot 0.5H_2O$: C, 60.68, H, 3.52, N, 3.08.

Found: C, 60.71, H, 3.91, N, 2.82.

EXAMPLE 31

5

4-(5-Chloro-2-hydroxyphenyl)-3-(4-acetamidophenyl)-6-(trifluoromethyl)-2(1H)-quinollnone (35e, $R^3 = CF_3$, $R^6 =$ 4-acetamidophenyl, $n = 0$)

10 mp 240-260°C;

1H (300 MHz, CD_3OD): δ 7.77 (1H, d, $J = 8.7$ Hz), 7.55 (1H, d, $J = 8.7$ Hz), 7.48-7.38 (m, 3H), 7.19-7.15 (m, 3H), 6.87-6.83 (m, 2H), 2.09 (s, 3H); MS m/e 472 (MH^+).

15

Anal. Calcd. for $C_{24}H_{16}ClF_3N_3O_3$: C, 56.59, H, 3.93, N, 5.50.

Found: C, 57.21, H, 3.73, N, 5.28.

EXAMPLE 32

20

4-(5-Chloro-2-hydroxyphenyl)-3-(4-aminophenyl)-6-(trifluoromethyl)-2(1H)-quinollnone (35f, $R^3 = CF_3$, $R^6 =$ 4-aminophenyl, $n = 0$)

25 mp 222-224°C;

1H (300 MHz, CD_3OD): δ 7.74 (1H, d, $J = 8.4$ Hz), 7.53 (1H, d, $J = 8.4$ Hz), 7.36 (s, 1H), 7.15 (dd, 1H, $J = 2.7$ and Hz), 6.99 (d, $J = 8.4$ Hz, 2H), 6.85-6.81 (m, 2H), 6.58 (1H, d, $J = 8.4$ Hz); MS m/e 446.8 (MH^+).

30

- 62 -

Anal. Calcd. for $C_{22}H_{14}ClF_3N_2O_3$: C, 59.14, H, 3.16, N, 6.27.

Found: C, 60.27, H, 3.52, N, 6.32.

EXAMPLE 33

5

4-(5-Chloro-2-hydroxyphenyl)-3-[2-(4-hydroxyphenyl)ethyl]-6-(trifluoromethyl)-2(1H)-quinolinone (35g, $R^3 = CF_3$, $R^8 =$ 4-hydroxyphenyl, $n = 2$)

10 mp 205-207°C;

1H (300 MHz, CD_3OD): δ 7.73-7.69 (1H, dd, $J = 1.8$ and 8.7 Hz), 7.52 (1H, d, $J = 8.7$ Hz), 7.38-7.35 (dd, 1H, $J = 2.7$ and 6.2 Hz), 7.23 (1H, s), 7.01 (d, 1H, $J = 8.7$ Hz), 6.78-6.60 (2H, dd, $J = 8.7$ and 7.3 Hz), 2.68 (m, 4H); MS m/e 459 (MH^+).

15

Anal. Calcd. for $C_{24}H_{17}ClF_3NO_3 \cdot 1.5H_2O$: C, 59.16, H, 4.11, N, 2.88.

Found: C, 58.71, H, 3.78, N, 2.86.

20

EXAMPLE 34

4-(5-Chloro-2-hydroxyphenyl)-3-methyl-6-(trifluoromethyl)-2(1H)-quinolinone (35h, $n = 0$, $R^8 = Me$, $R^3 = CF_3$)

25 MS m/z : 352 (MH^+);

IR (KBr) 3183, 1655, 1321, 1263, 1122 cm^{-1} ;

1H NMR ($DMSO-d_6$): δ 1.86 (3H, s), 7.04 (1H, d, $J=8.8$ Hz), 7.12 (1H, s), 7.22 (1H, d, $J=2.6$ Hz), 7.39 (1H, dd, $J=2.6, 8.7$ Hz), 7.52 (1H, d, $J=8.6$ Hz), 7.78 (1H, d, $J=8.7$ Hz), 9.92 (1H, s), 12.26 (1H, s);

30

- 63 -

Anal. Calcd. for $C_{17}H_{11}ClF_3NO_2 \cdot 0.5 H_2O$: C, 56.26; H, 3.33; N, 3.86.

Found: C, 56.57; H, 3.16; N, 3.81.

5

EXAMPLE 35

3-[4-(5-Chloro-2-hydroxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)quinolin-3-yl]acrylonitrile (36a, R = H, X = CN, R³ = CF₃)

10

To a cold suspension (0°C) of NaH (60% mineral oil, 33 mg, 0.82 mmol) in DMF (5 mL), diethyl cynomethylphosphonate (63 μ L, 0.39 mmol) was added dropwise. The reaction mixture was stirred at 0°C for 0.5 hours and a solution of 2-chloro-6,8-dihydro-6-hydroxy-11-(trifluoromethyl)-7H-[1]benzopyrano[3,4-c]quinolin-7-one prepared in Example 10, Step A (120 mg, 0.33 mmol) in DMF (5 mL) was added. The red reaction mixture was allowed to warm to room temperature and stirred for 2 hours. The reaction mixture was quenched with 1N HCl and then extracted with ethyl acetate. The organic layer was separated and washed with saturated NaHCO₃, water, brine and then dried (MgSO₄). Evaporation of the solvent gave a yellowish oil which was then purified by column chromatography (silica gel, 1:1 ethyl acetate/hexanes) to afford the title compound as a yellow solid (71 mg, 56%):

25 mp >265°C; MS m/e 389 (M-H)⁺.

Anal. Calcd. for $C_{19}H_{10}ClF_3N_2O_2$: C, 58.40; H, 2.58; N, 7.17.

Found: C, 58.16; H, 2.81; N, 6.87.

30 ¹H NMR (DMSO-d₆): δ 6.85 (d, J=16.4 Hz, 1H), 7.10 (d, J=8.8 Hz, 1H), 7.21(s, 1H), 7.23 (d, J=16.4 Hz, 1H), 7.30 (d, J=2.7 Hz, 1H), 7.49 (dd,

- 64 -

J=8.8 Hz, 2.7 Hz, 1H), 7.59 (d, J=8.6 Hz, 1H), 7.93 (dd, J=8.7 Hz, 1.7 Hz, 1H), 10.19 (s, 1H), 12.70 (s, 1H). IR (KBr, cm^{-1}): 3333, 2224, 1656, 1625, 1585, 1321, 1265, 1118, 1073.

5

EXAMPLE 36

3-[4-(5-Chloro-2-methoxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)quinolin-3-yl]acrylonitrile (36b, R = Me, X = CN, R³ = CF₃)

10

The title compound was prepared from the compound of the formula (5) prepared in Example 4 following the general procedure described in Example 35.

15 mp >250°C; MS m/e 403 (M-H)⁺.

¹H NMR (DMSO-d₆): δ 3.71 (3 H, s), 6.92 (1 H, d, J=16.3 Hz), 7.08 (2 H, m), 7.31 (1 H, s), 7.32 (1 H, d, J=16.3 Hz), 7.54 (2 H, m), 7.81 (1 H, dd, J=8.5 & 1.6 Hz), 12.41 (1 H, brd s). IR (KBr, cm^{-1}): 2216, 1665, 1321,

20 1127.

EXAMPLE 37(a)

4-[4-(5-Chloro-2-hydroxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)quinolin-3-yl]-3-buten-2-one (37a, R = H, X = Ac, R³ = CF₃)

25

The title compound was prepared from compound 13 prepared in Example 10, Step A by following the general procedure described in
30 Example 35.

- 65 -

mp 186-188°C. MS m/e: 406 (M-H).

IR (KBr, cm⁻¹): 3185, 1656, 1629, 1322, 1284, 1169, 1125, 1076

5 Anal. Calcd. for C₂₀H₁₃ClF₃NO₃: C, 58.91; H, 3.21; N, 3.43.

Found: C, 58.64; H, 3.05; N, 3.23.

EXAMPLE 37(b)

10 **4-[4-(5-Chloro-2-methoxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)quinolin-3-yl]-3-buten-2-one (37b, R = Me, X = Ac, R³ = CF₃)**

The title compound was prepared from compound 5 prepared in
15 Example 4 by following the general procedure described in Example 5.

mp 232-234°C. MS m/e: 422 (MH+).

IR (KBr, cm⁻¹): 2844, 1686, 1625, 1656, 1588, 1320.

20

Anal. Calcd. for C₂₁H₁₅ClF₃NO₃: C, 59.80; H, 3.58; N, 3.32.

Found: C, 59.60; H, 3.56; N, 3.22.

EXAMPLE 38

25

4-(5-Chloro-2-hydroxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-quinolinecarboxaldehyde oxime (38a, R = H, R³ = CF₃)

30

To a suspension of 2-chloro-6,8-dihydro-6-hydroxy-11-(trifluoromethyl)-7H-[1]benzopyrano[3,4-c]quinolin-7-one prepared in

- 66 -

Example 10, Step A (41 mg, 0.11 mmol) in THF (10 mL), hydroxylamine hydrochloride (9.3 mg, 0.13 mmol) and triethylamine (0.038 mL, 0.28 mmol) were added. The reaction mixture was stirred at room temperature overnight. The THF was evaporated and water was added. The light
5 yellow precipitate was collected and air dried to afford the title compound (36 mg, 85%): mp 195-198°C (dec.);

MS m/e 383 (MH⁺).

10 Anal. Calcd. for C₁₇H₁₀ClF₃N₂O₃: C, 53.35; H, 2.63; N, 7.32.
Found: C, 53.18; H, 4.55; N, 6.87.

¹H NMR (DMSO-d₆): δ 6.98 (d, J=8.8 Hz, 1H), 7.19-7.20 (m, 2H), 7.34
(m, 1H), 7.54 (d, J=8.6 Hz, 1H), 7.85 (m, 1H), 7.95 (s, 1H), 9.87 (s, br,
15 1H), 11.30 (s, 1H), 12.46 (s, br, 1H). IR (KBr, cm⁻¹): 3247, 1661, 1629,
1322, 1265, 1168, 1121, 1076.

EXAMPLE 39

20 **4-(5-Chloro-2-methoxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-quinolinecarboxaldehyde oxime (38b, R = Me, R³ = CF₃)**

A stirred suspension of the compound prepared in Example 4 (80
25 mg, 0.21 mmol), NH₂OH.HCl (18 mg, 0.25 mmol) and anhydrous NaOAc (20 mg, 0.25 mmol) in absolute ethanol (2 mL) was heated at reflux for 1 hour. The ethanol was rotary evaporated and the residue was partitioned between EtOAc and water. The EtOAc layer was separated and washed with water, brine and then dried (Na₂SO₄). Evaporation of EtOAc
30 followed by trituration of the crud product with ether gave the title compound as a white solid (56 mg): mp 255-258°C; IR (KBr, cm⁻¹) 3207,

- 67 -

1669, 1323, 1267, 1122;

¹H NMR (DMSO-d₆): δ 3.66 (3 H, s), 7.08 (1 H, s), 7.22 (1 H, d, J = 8.9 Hz), 7.28 (1 H, d, J = 2.6 Hz), 7.53 (2 H, dd, J = 8.9 and 2.8 Hz), 7.85 (1 H, dd, J = 8.7 and 1.7 Hz), 7.97 (1 H, s), 11.29 (1 H, s), 12.50 (1 H, brd s); MS 397 (MH⁺).

EXAMPLE 40

10 **4-(5-Chloro-2-hydroxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-quinolineacetic acid, methyl ester (30b, R^a = Me, R³ = CF₃)**

Step A: 2-Chloro-7,9-dihydro-12-(trifluoromethyl)-
15 [1]benzoxepino[4,5-c]quinolin-6,8-dione (39, R³ = CF₃)

A stirred mixture of the carboxylic acid of the formula (30a, R^a = H, R³ = CF₃) (1.20 g, 3.0 mmol) and a catalytic amount of *p*-TsOH was refluxed in toluene for 4 hours. The solvent was removed by rotary
20 evaporation. The residue was treated with a saturated solution of sodium bicarbonate and extracted with EtOAc. The organic extracts were dried over MgSO₄ and concentrated to give the title lactone compound (783 mg, 69%):

25 ¹H NMR (300 MHz, DMSO-d₆): δ 3.23 (1H, d, J = 13.3 Hz), 4.26 (1H, d, J = 13.3 Hz), 7.52 (1H, d, J = 8.8 Hz), 7.60 (1H, m), 7.76 (1H, dd, J = 8.8, 2.6 Hz), 7.87 (1H, d, J = 2.6 Hz), 7.92 (2H, m), 12.62 (1H, brd); MS *m/e* 380 (MH⁺).

30

- 68 -

Step B: 4-(5-Chloro-2-hydroxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-quinolineacetic acid, methyl ester (30b, $R^a = \text{Me}$, $R^b = \text{CF}_3$)

5 On attempted purification of the lactone 2-chloro-7,9-dihydro-12-(trifluoromethyl)-[1]benzoxepino[4,5-c]quinolin-6,8-dione prepared in Step A, by column chromatography on silica gel using a mixture of CH_2Cl_2 -MeOH as eluant gave the ester of the title compound.

10 ^1H NMR (300 MHz, $\text{DMSO}-d_6$): δ 3.17 (1H, d, $J = 16.4$ Hz), 3.47 (1H, d, $J = 16.4$ Hz), 3.53 (3H, s), 7.05 (1H, d, $J = 8.7$ Hz), 7.11 (1H, d, $J = 2.7$ Hz), 7.18 (1H, m), 7.42 (1H, dd, $J = 8.7, 2.7$ Hz), 7.55 (1H, d, $J = 8.5$ Hz), 7.84 (1H, dd, 8.7, 2.7 Hz), 10.0 (1H, s), 12.4 (1H, brd s); MS m/e 412 (MH^+).

15 **EXAMPLE 41**

4-(5-Chloro-2-hydroxyphenyl)-3-(2-hydroxy-2-methylpropyl)-6-(trifluoromethyl)-2(1H)-quinollnone (40, $R = R^1 = \text{Me}$, $R^3 = \text{CF}_3$)

20

A solution of methyl lithium (1M in THF, 1.6 mL, 1.6 mmol) was added to a cold (-78°C) stirred solution of 2-chloro-7,9-dihydro-12-(trifluoromethyl)-[1]benzoxepino[4,5-c]quinolin-6,8-dione prepared in Example 40, Step A (16 mg, 0.3 mmol) in anhydrous THF (3 mL) under nitrogen. After stirring for 1 hour at -78°C , the cold bath was removed and stirring was continued for 16 hours. The reaction was quenched with 1N HCl and then extracted with EtOAc. The crude product was purified by flash chromatography (silica gel, 19:1 $\text{CH}_2\text{Cl}_2/\text{MeOH}$) to afford the title compound (21 mg) as a big solid:

30

MS m/z : 412 (MH^+);

¹H NMR (DMSO-d₆): δ 0.95 (6H, s), 2.6 (2H, dd), 7.0 (1H, d), 7.1 (1H, s), 7.2 (1H, s), 7.39 (1H, d), 7.55 (1H, d), 7.8 (1H, d), 9.95 (1H, s), 12.5 (1H, s).

5

EXAMPLE 42

4-(5-Chloro-2-hydroxyphenyl)-1-methyl-3-(2-hydroxyethyl)-6-trifluoromethyl-2(1H)-quinolinone (43, R³ = CF₃)

10 **Step A:** 4-(5-Chloro-2-hydroxyphenyl)-3-(2-triisopropylsilyloxyethyl)-6-trifluoromethyl-2(1H)-quinolinone (41, R³ = CF₃)

Neat triisopropylsilyl chloride (0.293 mL, 1.37 mmol, 1.05
15 equivalent) was added to a stirred solution of the compound of Example 20 and imidazole (0.134 g, 1.97 mmol, 1.5 equivalent) in anhydrous DMF (10 mL). After 12 hours at room temperature, the resulting mixture was poured into aqueous 1 N HCl (50 mL). The aqueous layer was extracted with ethyl acetate (3 x 50 ml). The combined organic layers
20 were washed with brine (25 mL), dried (MgSO₄), filtered and concentrated in vacuo. The crude residue was purified using silica gel column chromatography (3:1-2:1 hexane/ethyl acetate) to afford 0.357 g of a clear viscous oil (50% yield). An analytical sample of the title compound was obtained by crystallization from ethyl acetate/hexane: mp
25 209-210°C.

¹H NMR (300 MHz, DMSO-d₆): δ 0.91 (21H, s), 2.58 (2H, m), 3.63 (2H, m), 7.04 (1H, d, J = 8.8), 7.06 (1H, s), 7.23 (1H, d, J = 2.6), 7.40 (1H, dd, J = 8.7, 2.3), 7.51 (1H, d, J = 8.6), 7.78 (1H, dd, J = 8.6, 1.7), 9.92 (1H, s),
30 12.26 (1H, s); MS m/e 540 (MH⁺).

- 70 -

Step B: 4-(5-Chloro-2-hydroxyphenyl)-1-methyl-3-(2-triisopropylsilyloxyethyl)-6-trifluoromethyl)-2(1H)-quinolinone (42, $R^3 = CF_3$)

5 *n*-Butyl lithium (0.593 mL, 0.950 mmol, 2.1 equivalent, 1.6 M/hexane) was added to a stirred solution of the quinolinone from Step A (0.244 g, 0.453 mmol) in THF at -78°C. After 15 minutes neat iodomethane was added and the resulting mixture was allowed to warm to room temperature. After stirring for 12 hours, the reaction was
10 quenched with aqueous 1 N HCl (10 mL) and poured into H₂O (50 mL). The aqueous layer was extracted with ethyl acetate (3 x 50 mL). The combined organic layers were washed with brine (25 mL), dried (MgSO₄), filtered and concentrated in vacuo. The crude residue was
15 purified using silica gel column chromatography (4:1-3:1 hexane/ethyl acetate) to afford 0.202 g of the title compound as a white solid (81% yield).

¹H NMR (300 MHz, DMSO-d₆): δ 0.92 (21H, s), 2.63 (2H, m), 3.66 (2H, t, *J* = 7.4), 3.75 (3H, s), 7.05 (1H, d, *J* = 8.8), 7.15 (1H, s), 7.24 (1H, d, *J* =
20 2.7), 7.41 (1H, dd, *J* = 8.7, 2.6), 7.78 (1H, d, *J* = 8.9), 7.91 (1H, dd, *J* = 9.0, 1.9), 9.95 (1H, s).

Step C: 4-(5-Chloro-2-hydroxyphenyl)-1-methyl-3-(2-hydroxyethyl)-6-trifluoromethyl)-2(1H)-quinolinone (43, $R^3 = CF_3$)

25 Tetrabutylammonium fluoride (0.380 mL, 0.308 mmol, 2 equivalent, 1.0 M/THF) was added to a stirred solution of the quinolinone from Step B (0.105 g, 0.190 mmol) in THF (10 mL). After 12 hours, the crude reaction mixture poured into H₂O (50 mL). The aqueous layer was
30 extracted with ethyl acetate (3 x 50 mL). The combined organic layers were washed with brine (25 mL), dried (MgSO₄), filtered and

- 71 -

concentrated in vacuo. The crude residue was purified using column chromatography (silica gel, 2.5% methanol/chloroform) to afford 74 mg of a white solid. The solid was recrystallized from ethyl acetate/hexane to provide 0.064 g (86% yield) of the title compound: mp 229-230°C.

5

^1H NMR (300 MHz, DMSO- d_6): δ 2.56 (2H, m), 3.40 (2H, m), 3.75 (3H, s), 4.57 (1 H, m), 7.05 (1H, d, J = 8.8), 7.14 (1H, s), 7.25 (1H, d, J = 2.6), 7.41 (1H, dd, J = 8.7, 2.7), 7.77 (1H, d, J = 8.8), 7.89 (1H, dd, J = 8.8, 2.0), 9.91 (1H, s); MS m/e 398 (MH^+);

10

Anal. Calcd. for $\text{C}_{19}\text{H}_{15}\text{ClF}_3\text{NO}_3$: C, 57.37; H, 3.80; N, 3.52.

Found: C, 57.69; H, 3.88; N, 3.28.

EXAMPLE 43

15

4-(5-Chloro-2-methoxyphenyl)-1-methyl-3-(2-hydroxyethyl)-6-trifluoromethyl)-2(1H)-quinolinone (44, $\text{R}^3 = \text{CF}_3$)

A mixture of the quinolinone compound prepared in Example 42, Step B (0.202 g, 0.365 mmol), dimethylsulfate (0.038 mL, 0.402 mmol, 1.1 equivalent), and potassium carbonate (0.056 g, 0.402 mmol, 1.1 equivalent) in acetone (15 mL) was heated at reflux for 3 hours. The resulting mixture was poured into H_2O (25 mL). The aqueous layer was extracted with ethyl acetate (3 x 50 mL). The combined organic layers were washed with brine (25 mL), dried (MgSO_4), filtered and concentrated in vacuo. The crude silylated quinolinone (0.228 g) was used in the subsequent reaction without purification or characterization.

Tetrabutylammonium fluoride (0.802 mL, 0.802 mmol, 2 equivalent, 1.0 M/THF) was added to a stirred solution of the silylated quinolinone (0.228 g, 0.401 mmol) in THF (10 mL). After 12 hours, the

30

- 72 -

crude reaction mixture poured into H₂O (50 mL). The aqueous layer was extracted with ethyl acetate (3 x 50 ml). The combined organic layers were washed with brine (25 mL), dried (MgSO₄), filtered and concentrated in vacuo. The crude residue was purified using column chromatography (silica gel, 1:1-1:2 hexane/ethyl acetate) to afford 0.148 g of a white solid. The solid was recrystallized from ethyl acetate/hexane to provide 0.139 g (84% yield) of the title compound: mp 175-176°C.

¹H NMR (300 MHz, DMSO-d₆): δ 2.48 (2H, m), 3.37 (2H, m), 3.67 (3H, s), 3.75 (3H, s), 4.57 (1 H, t, 5.4), 7.06 (1H, s), 7.29 (1H, d, J = 9.0), 7.35 (1H, d, J = 1.3), 7.61 (1H, dd, J = 8.8, 2.7), 7.77 (1H, d, J = 8.8), 7.89 (1H, dd, J = 8.8, 1.9); MS m/e 412 (MH⁺);

Anal. Calcd. for C₂₀H₁₇ClF₃NO₃: C, 58.33; H, 4.16; N, 3.40.
Found: C, 58.30; H, 4.07; N, 3.18.

EXAMPLE 44

4-(5-Chloro-2-hydroxyphenyl)-1-methyl-3-(2-hydroxyethyl)-6-trifluoromethyl)-2(1H)-quinolinone (43, R³ = CF₃)

Step A: 4-(5-Chloro-2-hydroxyphenyl)-3-(2-trisopropylsilyloxyethyl)-6-trifluoromethyl)-2(1H)-quinolinone (41, R³ = CF₃)

Neat triisopropylsilyl chloride (0.293 mL, 1.37 mmol, 1.05 equivalent) was added to a stirred solution of the quinolone **31a** prepared in Example 20 and imidazole (0.134 g, 1.97 mmol, 1.5 equivalent) in anhydrous DMF (10 mL). After 12 hours at room temperature, the resulting mixture was poured into aqueous 1 N HCl (50 mL). The aqueous layer was extracted with ethyl acetate (3 x 50 ml).

- 73 -

The combined organic layers were washed with brine (25 mL), dried (MgSO₄), filtered and concentrated in vacuo. The crude residue was purified using silica gel column chromatography (3:1-2:1 hexane/ethyl acetate) to afford 0.357 g of a clear viscous oil (50% yield). An analytical sample was obtained by crystallization from ethyl acetate/hexane: mp 209-210°C.

¹H NMR (300 MHz, DMSO-d₆): δ 0.91 (21H, s), 2.58 (2H, m), 3.63 (2H, m), 7.04 (1H, d, *J* = 8.8), 7.06 (1H, s), 7.23 (1H, d, *J* = 2.6), 7.40 (1H, dd, *J* = 8.7, 2.3), 7.51 (1H, d, *J* = 8.6), 7.78 (1H, dd, *J* = 8.6, 1.7), 9.92 (1H, s), 12.26 (1H, s); MS *m/e* 540 (MH⁺).

Step B: 4-(5-Chloro-2-hydroxyphenyl)-1-methyl-3-(2-triisopropylsilyloxyethyl)-6-trifluoromethyl)-2(1H)-quinolinone (42, R³ = CF₃)

n-Butyl lithium (0.593 mL, 0.950 mmol, 2.1 equivalent, 1.6 M/hexane) was added to a stirred solution of the quinolinone 41 (0.244 g, 0.453 mmol) prepared in Step A, in THF at -78°C. After 15 minutes, neat iodomethane was added and the resulting mixture was allowed to warm to room temperature. After stirring for 12 hours, the reaction was quenched with aqueous 1 N HCl (10 mL) and poured into H₂O (50 mL). The aqueous layer was extracted with ethyl acetate (3 x 50 mL). The combined organic layers were washed with brine (25 mL), dried (MgSO₄), filtered and concentrated in vacuo. The crude residue was purified using silica gel column chromatography (4:1-3:1 hexane/ethyl acetate) to afford 0.202 g of the title compound as a white solid (81% yield).

¹H NMR (300 MHz, DMSO-d₆): δ 0.92 (21H, s), 2.63 (2H, m), 3.66 (2H, t, *J* = 7.4), 3.75 (3H, s), 7.05 (1H, d, *J* = 8.8), 7.15 (1H, s), 7.24 (1H, d, *J* =

- 74 -

2.7), 7.41 (1H, dd, $J = 8.7, 2.6$), 7.78 (1H, d, $J = 8.9$), 7.91 (1H, dd, $J = 9.0, 1.9$), 9.95 (1H, s).

Step C: 4-(5-Chloro-2-hydroxyphenyl)-1-methyl-3-(2-hydroxyethyl)-
5 6-trifluoromethyl)-2(1H)-quinolinone (43, $R^3 = CF_3$)

Tetrabutylammonium fluoride (0.380 mL, 0.308 mmol, 2
equivalent, 1.0 M/THF) was added to a stirred solution of the quinolinone
42 (0.105 g, 0.190 mmol) prepared in Step B, in THF (10 mL). After 12
10 hours, the crude reaction mixture poured into H₂O (50 mL). The
aqueous layer was extracted with ethyl acetate (3 x 50 ml). The
combined organic layers were washed with brine (25 mL), dried
(MgSO₄), filtered and concentrated in vacuo. The crude residue was
purified using column chromatography (silica gel, 2.5%
15 methanol/chloroform) to afford 74 mg of a white solid. The solid was
recrystallized from ethyl acetate/hexane to provide 0.064 g (86% yield) of
the title compound: mp 229-230°C.

¹H NMR (300 MHz, DMSO-d₆): δ 2.56 (2H, m), 3.40 (2H, m), 3.75 (3H,
20 s), 4.57 (1 H, m), 7.05 (1H, d, $J = 8.8$), 7.14 (1H, s), 7.25 (1H, d, $J = 2.6$),
7.41 (1H, dd, $J = 8.7, 2.7$), 7.77 (1H, d, $J = 8.8$), 7.89 (1H, dd, $J = 8.8, 2.0$),
9.91 (1H, s); MS m/e 398 (MH⁺);

Anal. Calcd. for C₁₉H₁₅ClF₃NO₃: C, 57.37; H, 3.80; N, 3.52.
25 Found: C, 57.69; H, 3.88; N, 3.28.

EXAMPLE 45

4-(5-Chloro-2-methoxyphenyl)-1-methyl-3-(2-hydroxyethyl)-6-
30 trifluoromethyl)-2(1H)-quinolinone (44, $R^3 = CF_3$)

- 75 -

A mixture of the quinolinone 42 prepared in Example 44, Step B (0.202 g, 0.365 mmol), dimethylsulfate (0.038 mL, 0.402 mmol, 1.1 equivalent), and potassium carbonate (0.056 g, 0.402 mmol, 1.1 equivalent) in acetone (15 mL) was heated at reflux for 3 hours. The resulting mixture was poured into H₂O (25 mL). The aqueous layer was extracted with ethyl acetate (3 x 50 mL). The combined organic layers were washed with brine (25 mL), dried (MgSO₄), filtered and concentrated in vacuo. The crude quinolinone (0.228 g) was used in the subsequent reaction without purification or characterization.

10

Tetrabutylammonium fluoride (0.802 mL, 0.802 mmol, 2 equivalent, 1.0 M/THF) was added to a stirred solution of the quinolinone prepared above (0.228 g, 0.401 mmol) in THF (10 mL). After 12 hours, the crude reaction mixture was poured into H₂O (50 mL). The aqueous layer was extracted with ethyl acetate (3 x 50 mL). The combined organic layers were washed with brine (25 mL), dried (MgSO₄), filtered and concentrated in vacuo. The crude residue was purified using column chromatography (silica gel, 1:1-1:2 hexane/ethyl acetate) to afford 0.148 g of a white solid. The solid was recrystallized from ethyl acetate/hexane to provide 0.139 g (84% yield) of the title compound: mp 175-176°C.

¹H NMR (300 MHz, DMSO-d₆): δ 2.48 (2H, m), 3.37 (2H, m), 3.67 (3H, s), 3.75 (3H, s), 4.57 (1 H, t, 5.4), 7.06 (1H, s), 7.29 (1H, d, J = 9.0), 7.35 (1H, d, J = 1.3), 7.61 (1H, dd, J = 8.8, 2.7), 7.77 (1H, d, J = 8.8), 7.89 (1H, dd, J = 8.8, 1.9); MS m/e 412 (MH⁺);

25

Anal. Calcd. for C₂₀H₁₇ClF₃NO₃: C, 58.33; H, 4.16; N, 3.40.

Found: C, 58.30; H, 4.07; N, 3.18.

- 76 -

EXAMPLE 46**4-(5-Chloro-2-methoxyphenyl)-3-methyl-6-trifluoromethyl-2(1H)-quinolinone**

5

A solution of compound 33 ($R^3 = CF_3$, $R^8 = H$, $n = 1$) (5.63 mmol), 33% HBr in AcOH (38.3 mmol) and 10 mL of AcOH was heated to 75°C for 3 hours. The solution was cooled to room temperature and quenched with H₂O (50 mL) and then agitated for 12 hours. The precipitate was
10 filtered, washed with H₂O, and dried *in vacuo*. The pale brown solid was recrystallized from ethyl acetate/hexane. The title compound was isolated as a white solid (0.550 g, 27% yield).

¹H NMR (300 MHz, CDCl₃): δ 2.04 (s, 3H), 3.72 (s, 3H), 7.03 (d, 1H, $J =$
15 9.0 Hz), 7.12 (s, 1H), 7.44 (m, 3H), 7.65 (d, 1H, $J = 8.4$ Hz), 10.93 (br s, 1H); MS m/e 368 (MH⁺);

Anal. Calcd. for C₁₈H₁₃ClF₃NO₂ • 0.33 H₂O: C, 58.79; H, 3.56; N, 3.81.

Found: C, 58.89; H, 3.82; N, 3.53.

20

EXAMPLE 47**4-(5-Chloro-2-hydroxyphenyl)-3-(2-hydroxyethyl)-6-(trifluoromethyl)-2(1H)-quinolinone (31a, $R^3 = CF_3$, $n = 2$)**

25

Step A: 3-(2-Hydroxyethyl)-4-hydroxy-6-chlorocoumarin (45)

To a solution of γ -butyrolactone (15.5 g, 178.0 mmol) in THF (100 mL) at -78°C was added a 1.0 M THF solution of LiHMDS (356 mL, 356 mmol),
30 and the resulting mixture stirred at -78°C for 1.5 hours. A solution of 5-

- 77 -

chlorosalicylic methyl ester (16.6 g, 98% purity, 89.0 mmol) in THF (95 mL) was added. After stirring for 1 hour at 0°C, the mixture was warmed to room temperature overnight to ensure complete reaction. After cooling to 0°C, conc. HCl (12 N, 150 mL) was slowly added to bring the pH to 1. The reaction solution was stirred until HPLC analysis indicated the absence of the keto-ester intermediate. To the mixture was added 400 mL CH₂Cl₂ and 300 mL H₂O; the organic phase was separated and the aqueous layer was extracted with CH₂Cl₂ (100 mL). The organic layers were combined and dried over anhydrous Na₂SO₄, and the solvent was removed under reduced pressure to give a solid. Heptane (165 mL) was added to a solution of the solid in THF (290 mL) to crystallize the product. After cooling to 0-5°C for about 3 hours, the product was isolated by filtration and washed with heptane. After drying in vacuo, a total of 13.9 g (66% yield) of the title compound as off-white crystals was obtained. m.p. 185-186°C; MS m/z 240;

¹H NMR (DMSO, 300 MHz) δ 7.84 (d, 1H, *J* = 2.4 Hz), 7.61 (dd, 1H, *J* = 2.4, 8.8 Hz), 7.38 (d, 1H, *J* = 8.8 Hz), 3.56 (t, 2H, *J* = 6.6 Hz), 2.73 (t, 2H, *J* = 6.6 Hz); ¹³C NMR (DMSO, 75 MHz) δ 162.6, 159.9, 150.5, 131.4, 127.9, 122.4, 118.2, 117.8, 103.2, 59.4, 27.6; IR (cm⁻¹) 3247.2, 2945.1, 2458.6, 1664.9, 1623.9, 1572.7, 1311.5, 1378.1, 1070.8, 825.0.

Step B: 2,3-Dihydro-8-chloro-4H-furobenzopyran-4-one (46)

To a solution of 3-(2-hydroxyethyl)-4-hydroxy-6-chlorocoumarin (45) (8 g, 33.3 mmol) in toluene (360 mL) at room temperature was added *p*-TSA (0.95 g, 5.0 mmol), and the resulting solution was refluxed with the removal of water using a Dean-Stark condenser. The reaction mixture was cooled to room temperature and washed with saturated sodium bicarbonate solution twice. Toluene was removed by atmospheric distillation to a final volume of 32 mL. After cooling to 70°C, the product

- 78 -

started to crystallize. The crystal slurry was held between 55-65°C for 30 minutes, followed by cooling to 0-5°C. The product was isolated by filtration, washed with cold toluene, and dried in vacuo. A total of 5.5 g (74% yield) of the title compound as off-white crystals was obtained.

5 m.p. 144-146°C; MS m/z 223 (M+H);

¹H NMR (CDCl₃, 300 MHz) δ 7.58 (d, 1H, J = 2.5 Hz), 7.49 (dd, 1H, J = 2.3, 8.8 Hz), 7.30 (d, 1H, J = 8.9 Hz), 4.90 (t, 2H, J = 9.3 Hz), 3.21 (t, 2H, J = 9.5 Hz); ¹³C NMR (CDCl₃, 75 MHz) δ 166.4, 160.3, 153.4, 132.6, 129.6,
10 122.4, 118.6, 113.8, 103.6, 74.9, 27.1; IR (cm⁻¹) 3073.1, 2975.8, 1721.2, 1644.4, 1490.8, 1403.7, 1270.6, 1111.8, 1040.1.

Step C: 4-(4'-Trifluoromethylphenylcarboxamide)-5-(2-hydroxy-5-chloro)-2,3-dihydrofuran (47)

15

To a solution of 2,3-dihydro-8-chloro-4H-furobenzopyran-4-one (46) (1.02 g, 4.58 mmol) and 4-(trifluoromethyl)aniline (0.74 g, 4.58 mmol) in THF (50 mL) at -15°C was added LiHMDS (10.5 mL, 10.5 mmol, 1.0M solution in THF). The clear, red solution was stirred at -15°C until HPLC
20 analysis indicated <1% of compound (46) remained (approximately 30 minutes). The reaction mixture was quenched by the addition of an aqueous solution of NaH₂PO₄ (50 mL, 10 wt% in H₂O). After the addition of MTBE (25 mL), the layers were separated and the rich organic phase washed successively with NaH₂PO₄ (50 mL, 10 wt % in H₂O) and
25 saturated brine solution. After drying over Na₂SO₄, the solution was concentrated to give the title compound as a clear, orange oil (1.76 g, 100% yield) which crystallized upon refrigeration. Addition of dichloromethane (20 mL) gave white crystals, which were isolated by filtration, washed with dichloromethane (10 mL) and dried to give 1.6

- 79 -

grams of the title compound (90% yield). m.p 180 –180.5°C; MS m/z 384 (M+H);

¹H NMR (DMSO, 300 MHz) δ 9.76 (s, 1H), 9.34 (s, 1H), 7.76 (d, 2H, J = 8.5 Hz), 7.60 (d, 2H, J = 8.7 Hz), 7.26 (s, 1H), 7.24 (dd, 1 H, J = 2.2, 7.0 Hz), 6.83 (dd, 1H, J = 2.4, 7.1), 4.52 (t, 2H, J = 9.6 Hz), 3.16 (t, 2H, J = 9.6 Hz); ¹³C NMR (DMSO, 75 MHz) δ 165.5, 159.7, 155.9, 144.7, 132.0, 131.3, 127.3, 123.7, 121.7, 121.2, 119.5, 110.1, 71.5, 32.9; IR (cm⁻¹) 3303.6, 2950.2, 1654.6, 1608.5, 1531.7, 1408.8, 1326.9, 1116.9, 1065.7, 840.4.


Step D: 4-(5-Chloro-2-hydroxyphenyl)-3-(2-hydroxyethyl)-6-(trifluoromethyl)-2(1H)-quinolinone (31a, R³ = CF₃, n = 2)

A solution of 4-(4'-trifluoromethylphenyl)carboxamide)-5-(2-hydroxy-5-chloro)-2,3-dihydrofuran (47) prepared in Step C (1.76 g, 4.58 mmol) in MeOH (500 mL) was purged with nitrogen and irradiated with a 450 W Hanovia lamp at 30-40°C until HPLC analysis indicated < 1% of compound (47) remained. The MeOH was then concentrated in vacuo, and the resulting oil dissolved in dichloromethane (50 mL). Crystals formed after stirring for one hour at room temperature. After cooling the slurry to 0 °C, the crystals were isolated by filtration and dried. A total of 0.54 g (30% yield) of the title compound was obtained as a crystalline solid with an HPLC purity of 97 area % and had physical characterizing data which was identical to the compound of Example 20.

- 80 -

What is claimed:

1. A compound of the formula



5

wherein

R and R¹ each are independently hydrogen or methyl;

R², R³

and R⁴ each are independently hydrogen, halogen, nitro or trifluoromethyl, provided R², R³, and R⁴ are not all hydrogen;

10

R⁵ is bromo, chloro or nitro;

R⁶ is hydrogen or fluoro;

n is an integer from 0 to 6;


15 m is an integer of 0 or 1; and

R⁷ is CH₃, -CRR¹OH, -CHO, -C=NOH, -COCH₃ or aryl optionally substituted by one or two substituents selected from the group consisting of halogen, hydroxy, methoxy, amino, acetamino and trifluoromethyl;

20 or a nontoxic pharmaceutically acceptable salt thereof.

2. The compound of claim 1 having the formula

- 81 -



wherein

R is hydrogen or methyl;

5 R², R³

and R⁴ each are independently hydrogen, halogen, nitro or trifluoromethyl, provided R², R³, and R⁴ are not all hydrogen;

R⁵ is chloro;

10 R⁶ is hydrogen or fluoro;

n is an integer from 0 to 3;

m is an integer of 0 or 1; and

R⁷ is -CH₂OH, -CHO, -C=NOH, or aryl optionally substituted by one or two substituents selected from the group consisting of halogen, hydroxy, methoxy, amino, acetamino and trifluoromethyl;

15

or a nontoxic pharmaceutically acceptable salt thereof.

3. The compound of the claim 2 wherein R³ is trifluoromethyl, R² and R⁴ are hydrogen, and R⁷ is -CH₂OH; or a nontoxic pharmaceutically acceptable salt thereof.

20

4. The compound of claim 2 wherein R³ is trifluoromethyl, R² and R⁴ are hydrogen, and R⁷ is aryl optionally substituted by one or two substituents selected from the group consisting of halogen, hydroxy, methoxy, amino, acetamino and trifluoromethyl; or a nontoxic pharmaceutically acceptable salt thereof.

25

5. The compound of claim 1 selected from the group consisting of:

- 4-(5-chloro-2-methoxyphenyl)-3-(hydroxymethyl)-6-(trifluoromethyl)-
2(1H)-quinolinone;
- 5 4-(5-chloro-2-methoxyphenyl)-3-(hydroxymethyl)-7-(trifluoromethyl)-
2(1H)-quinolinone;
- 4-(5-chloro-2-methoxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-
quinolinecarboxaldehyde;
- 4-(5-chloro-2-methoxyphenyl)-3-(3-hydroxy-1-propenyl)-6-
10 (trifluoromethyl)-2(1H)-quinolinone;
- 4-(5-chloro-2-methoxyphenyl)-3-(3-hydroxypropyl)-6-(trifluoromethyl)-
2(1H)-quinolinone;
- 4-(5-chloro-2-hydroxyphenyl)-3-(hydroxymethyl)-6-(trifluoromethyl)-
2(1H)-quinolinone;
- 15 4-(5-chloro-2-hydroxyphenyl)-3-(3-hydroxy-1-propenyl)-6-
(trifluoromethyl)-2(1H)-quinolinone;
- 4-(5-chloro-2-hydroxyphenyl)-3-(3-hydroxypropyl)-6-(trifluoromethyl)-
2(1H)-quinolinone;
- (E)-4-(5-chloro-2-hydroxyphenyl)-3-(2-fluoro-3-hydroxy-1-propenyl)-6-
20 trifluoromethyl)-2(1H)-quinolinone;
- (Z)-4-(5-chloro-2-hydroxyphenyl)-3-(2-fluoro-3-hydroxy-1-propenyl)-6-
trifluoromethyl)-2(1H)-quinolinone;
- (E)-4-(5-chloro-2-methoxyphenyl)-3-(2-fluoro-3-hydroxy-1-propenyl)-6-
(trifluoromethyl)-2(1H)-quinolinone;
- 25 (Z)-4-(5-chloro-2-methoxyphenyl)-3-(2-fluoro-3-hydroxy-1-propenyl)-6-
(trifluoromethyl)-2(1H)-quinolinone;
- 4-(5-chloro-2-hydroxyphenyl)-3-(2-hydroxyethyl)-6-(trifluoromethyl)-
2(1H)-quinolinone;
- 4-(5-chloro-2-methoxyphenyl)-3-(2-hydroxyethyl)-6-trifluoromethyl-
30 2(1H)-quinolinone;
- 4-(5-chloro-2-methoxyphenyl)-3-(4-methoxyphenyl)-6-(trifluoromethyl)-
2(1H)-quinolinone;

- 83 -

- 4-(5-chloro-2-methoxyphenyl)-3-[(4-methoxyphenyl)methyl]-6-(trifluoromethyl)-2(1H)-quinolinone;
4-(5-chloro-2-methoxyphenyl)-3-(4-nitrophenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
5 4-(5-chloro-2-methoxyphenyl)-3-(4-aminophenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
4-(5-chloro-2-hydroxyphenyl)-3-(3,4-dimethoxyphenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
4-(5-chloro-2-hydroxyphenyl)-3-(2,4-dihydroxyphenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
10 4-(5-chloro-2-hydroxyphenyl)-3-(4-hydroxyphenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
4-(5-chloro-2-hydroxyphenyl)-3-[(4-hydroxyphenyl)methyl]-6-(trifluoromethyl)-2(1H)-quinolinone;
15 4-(5-chloro-2-hydroxyphenyl)-3-(4-acetamidophenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
4-(5-chloro-2-hydroxyphenyl)-3-(4-aminophenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
4-(5-chloro-2-hydroxyphenyl)-3-[2-(4-hydroxyphenyl)ethyl]-6-(trifluoromethyl)-2(1H)-quinolinone;
20 4-(5-chloro-2-hydroxyphenyl)-3-methyl-6-(trifluoromethyl)-2(1H)-quinolinone;
4-[4-(5-chloro-2-hydroxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)quinolin-3-yl]-3-buten-2-one;
25 4-(5-chloro-2-hydroxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-quinolinecarboxaldehyde oxime;
4-(5-chloro-2-methoxyphenyl)-1,2-dihydro-2-oxo-6-(trifluoromethyl)-3-quinolinecarboxaldehyde oxime; and
4-(5-chloro-2-hydroxyphenyl)-3-(2-hydroxy-2-methylpropyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
30 or a nontoxic pharmaceutically acceptable salt thereof.

- 84 -

6. The compound of claim 5 selected from the group consisting of:

- 4-(5-chloro-2-methoxyphenyl)-3-(3-hydroxy-1-propenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
- 5 4-(5-chloro-2-methoxyphenyl)-3-(3-hydroxypropyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
- 4-(5-chloro-2-hydroxyphenyl)-3-(hydroxymethyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
- 4-(5-chloro-2-hydroxyphenyl)-3-(3-hydroxy-1-propenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
- 10 4-(5-chloro-2-hydroxyphenyl)-3-(3-hydroxypropyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
- 4-(5-chloro-2-hydroxyphenyl)-3-(2-hydroxyethyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
- 15 4-(5-chloro-2-methoxyphenyl)-3-(2-hydroxyethyl)-6-trifluoromethyl-2(1H)-quinolinone;
- 4-(5-chloro-2-hydroxyphenyl)-3-(4-hydroxyphenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
- 4-(5-chloro-2-hydroxyphenyl)-3-[(4-hydroxyphenyl)methyl]-6-(trifluoromethyl)-2(1H)-quinolinone; and
- 20 4-(5-chloro-2-hydroxyphenyl)-3-(4-aminophenyl)-6-(trifluoromethyl)-2(1H)-quinolinone;
- or a nontoxic pharmaceutically acceptable salt thereof.

25 7. The compound of claim 1 which is 4-(5-chloro-2-hydroxyphenyl)-3-(2-hydroxyethyl)-6-(trifluoromethyl)-2(1H)-quinolinone.

8. The compound of claim 1 which is 4-(5-chloro-2-methoxyphenyl)-3-(2-hydroxyethyl)-6-trifluoromethyl-2(1H)-quinolinone.

30

9. A pharmaceutical composition for the treatment of disorders responsive to openers of the large conductance calcium-activated

- 85 -

potassium channels comprising a therapeutically effective amount of a compound as defined in claim 1 in association with a pharmaceutically acceptable carrier or diluent.

5 10. A method for the treatment of disorders responsive to opening of the large conductance calcium-activated potassium channels in a mammal in need thereof, which comprises administering to said mammal a therapeutically effective amount of a compound as defined in claim 1.

10

11. The method of claim 10 wherein said disorder is ischemia, stroke, convulsions, epilepsy, asthma, irritable bowel syndrome, migraine, traumatic brain injury, spinal cord injury, sexual dysfunction and urinary incontinence.

15

12. The method of claim 10 wherein said disorder is male erectile dysfunction.

20

13. The method of claim 11 wherein said disorder is stroke or traumatic brain injury.

14. The method of claim 11 wherein said disorder is sexual dysfunction.

25

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/28428

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07D 215/227
US CL : 514/312; 546/157

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
U.S. : 514/312; 546/157

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
EAST, Cas Online, USPatFull, Registry

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Database CA on STN, Chem Abstracts, (Columbus OH, USA) No. 129:41086, Sit, Sing-Yuen et al. '4-Aryl-3-hydroxyquinoline-2-one derivatives as ion channel modulator', abstract WO 9823273, June 1998.	1-14
A,P	Database USPATFULL on STN, US Patents Full Text, (Columbus, OH, USA) No. 1999:132843, Hewawasam, Piyasena et al., '2-aryl-3-aminoquinoline-2-one derivatives as potassium channel modulator', abstract US Patent 5972961, October 1999.	1-14
A,P	Database USPATFULL on STN, US Patents Full Text, (Columbus OH, USA) No. 1999:78740, Si, Sint-Yuen et al., '4-aryl-3-hydroxy-quinolin-2-one derivatives as ion channel modulators', abstract US Patent 5922735, July 1999.	1-14
A,P	Database CA on STN, Chem Abstracts, (Columbus OH, USA) No. 130:252386, Sit, Sing-Yuen et al. '4-aryl-3-hydroxyquinolin-2-one derivatives as ion channel modulators', abstract US Patent 5892045, April 1999.	1-14
A,P	Database CA on STN, Chem Abstracts, (Columbus OH, USA) No. 130:223174, Hewawasam, Piyasena et al., 'Preparation of 4-aryl-3-aminoquinoline-2-ones as potassium channel modulators', abstract WO 9909983, March 1999.	1-14



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	"&" document member of the same patent family

Date of the actual completion of the international search

04 April 2000 (04.04.2000)

Date of mailing of the international search report

26 APR 2000

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

D. Margaret Seaman
D. Margaret Seaman

Telephone No. (703) 308-0196